

Discrete and smooth orthogonal systems: C^∞ -approximation

A.I. Bobenko ^{*} D. Matthes [†] Yu.B. Suris [†]

Institut für Mathematik, Technische Universität Berlin,
Str. des 17. Juni 136, 10623 Berlin, Germany.

E-Mails: `bobenko@math.tu-berlin.de`, `matthes@math.tu-berlin.de`,
`suris@sfb288.math.tu-berlin.de`

1 Introduction and main results

Triply orthogonal coordinate systems have attracted the attention of mathematicians and physicists for almost two hundred years now. Particular examples of them were already used by Leibniz and Euler to evaluate multiple integrals in canonical coordinates, and later Lamé and Jacobi carried out calculations in analytical mechanics with the help of the famous elliptic coordinates. The first general results on the geometry of triply orthogonal systems date back to the 19th century – like the famous theorem of Dupin, saying that coordinate surfaces intersect along curvature lines. The Lamé equations which analytically describe the triply orthogonal systems were studied in detail by Bianchi [Bi2] and Darboux [Da].

Recently orthogonal systems came back into the focus of interest in mathematical physics as an example of an integrable system. Zakharov [Z] has shown how the Lamé equations can be solved by the $\bar{\partial}$ -method and constructed a variety of explicit solutions with the help of the dressing method. Algebraic-geometric solutions of the Lamé equations were constructed by Krichever [K]. The recent interest to the orthogonal coordinate systems is in particular motivated by their applications to the theory of the associativity equations [Du].

^{*}Partially supported by the SFB 288 “Differential Geometry and Quantum Physics” and the DFG Research Center “Mathematics for Key Technologies” (FZT 86) in Berlin.

[†]Supported by the SFB 288 “Differential Geometry and Quantum Physics”.




Figure 1: An elementary hexahedron of a discrete orthogonal system

The question of proper discretization of the classical models in differential geometry became recently a subject of intensive study (see, in particular, [BP, DS2, KS]). Indeed one can suggest various models in discrete geometry which have the same continuous limit and nevertheless have quite different properties. For a great variety of geometric problems described by integrable equations it was found that *integrable discretizations* (i.e. the discretizations preserving the integrability of the underlying nonlinear system) preserve the characteristic geometric properties of the problem. Moreover, it turns out that this discretization can be described in pure geometric terms.

Such a discretization of the triply-orthogonal coordinate systems was first suggested by one of the authors [Bo]. It is based on the classical Dupin theorem. Since the circular nets are known [MPS, N] to correspond to the curvature line parametrized surfaces, it is natural to define discrete triply orthogonal systems as maps from $\mathbb{Z}^3$ (or a subset thereof) to $\mathbb{R}^3$ with all elementary hexahedrons lying on spheres. The neighboring spheres intersect along circles which build the coordinate discrete curvature line nets (see Fig.3). Doliwa and Santini [CDS] made the next crucial step in the development of the theory. They considered discrete orthogonal systems as a reduction of discrete conjugated systems [DS1], generalized them to arbitrary dimension and proved their geometric integrability based on the classical Miguel theorem [Be]. The latter claims that provided the seven (black) points lie on circles as shown in Fig.1, the three dashed circles intersect in a common (white) point. This implies that a discrete triply-orthogonal system is uniquely determined by its three coordinate circular nets (as shown in Fig.3). This is a well-posed initial value problem for triply orthogonal nets.

Later on discrete orthogonal systems were treated by analytic methods of the theory of solitons. Algebro-geometric solutions [AKV] as well as the $\bar{\partial}$ -method [DMS] were discretized preserving all the symmetries of the smooth




Figure 2: Initial data for a smooth triply orthogonal system.




Figure 3: Initial data for a discrete triply orthogonal system.

case.

There is a common belief that the smooth theory can be obtained as a limit of the corresponding discrete one, which can be treated also in this case as a practical geometric numerical scheme for computation of smooth surfaces. Moreover, this discrete master theory has an important advantage: surfaces and their transformations appear exactly in the same way as sublattices of multidimensional lattices. This description immediately provides the statements about the permutability of the corresponding (Bäcklund-Darboux) transformations, – an important property which is nontrivial to see and sometimes even difficult to check in the smooth theory.

Until recently there were no rigorous mathematical statements supporting the observation about the classical limit of the discrete theories. The first step in closing this gap was made in our paper [BMS] where general convergence results were proven for a geometric numerical scheme for a class of nonlinear hyperbolic equations. The first geometric example covered by this theory was smooth and discrete surfaces with constant negative Gaussian curvature described by the sine-Gordon equation.

In the present paper we formulate initial value problems and prove the corresponding convergence results for conjugate and orthogonal coordinate systems, which are described by three-dimensional partial differential/difference equations. One should note here that despite the known geometric discretization, a convenient analytic description of discrete orthogonal systems converging to the classical description of the smooth case was missing. In particular this was the problem with discrete Lamé equations (cf. [AKV]).

The Clifford algebra description of smooth and discrete orthogonal systems suggested in [BH] turned out to be optimal for our purposes. This description provides us with the proper discrete analogs of the continuous quantities and their equations.

In particular we introduce the discrete Lamé system which is the starting point for further analytical investigations. We write these equations in the form of a discrete hyperbolic system and formulate the corresponding Cauchy problem. As a by-product, we rewrite the classical Lamé system in the hyperbolic form and pose a Cauchy problem for classical orthogonal coordinate systems, which seems to be new.

The central theorems of this paper are on the convergence of discrete orthogonal coordinate systems to a continuous one. In particular we prove the following statement.

Theorem 1 *Assume three umbilic free immersed surfaces $\mathcal{F}_1$, $\mathcal{F}_2$ and $\mathcal{F}_3$ in $\mathbb{R}^3$ intersect pairwise along their curvature lines, $X_1 = \mathcal{F}_2 \cap \mathcal{F}_3$ etc. Then:*

1. *On a sufficiently small box $B = [0, r]^3 \subset \mathbb{R}^3$, there is an orthogonal coordinate system $x : B \rightarrow \mathbb{R}^3$, which has $\mathcal{F}_1$, $\mathcal{F}_2$ and $\mathcal{F}_3$ locally as images of the coordinate planes, with each curve X_i being parametrized locally over the i -th axis by arc-length. The orthogonal coordinate system x is uniquely determined by these properties.*
2. *Denote by $B^\epsilon = B \cap (\epsilon\mathbb{Z})^3$ the grid of mesh size ϵ inside B . One can define a family $\{x^\epsilon : B^\epsilon \rightarrow \mathbb{R}^3\}$ of discrete orthogonal systems, parametrized by $\epsilon > 0$, such that the approximation error decays as*

$$\sup_{\xi \in B^\epsilon} |x^\epsilon(\xi) - x(\xi)| < C\epsilon.$$

The convergence is C^∞ : all partial difference quotients of x^ϵ converge with the same rate to the respective partial derivatives of x .

We also answer the question of *how to construct* the approximating family $\{x^\epsilon\}$: One solves the discrete Lamé system with discrete Cauchy data determined by $\mathcal{F}_1$, $\mathcal{F}_2$ and $\mathcal{F}_3$. The solution x^ϵ is then calculated in a process that consists of $\mathcal{O}(\epsilon^{-3})$ -many evaluations of the expressions in the discrete Lamé equations. The scheme can be – and has been – easily implemented on a computer, it needs a precision of ϵ^2 for the calculations and is robust against C^∞ -deviations of order ϵ in the initial data.

Actually, the results presented here go further:

- We do not restrict ourselves to the case of three dimensions, but allow arbitrary dimensions, possibly different for domain and target space.

In particular, we prove the corresponding approximation results for curvature line parametrized surfaces.

- One can keep some of the directions of the orthogonal system discrete while the others become continuous. This way, the classical Ribaucour transformation is obtained very naturally. The approximation result holds simultaneously for the system and its Ribaucour transforms.
- The results about the permutability of the Ribaucour transformations are elementary in the discrete setup. Our approximation theorem implies the corresponding claim for classical orthogonal systems (Theorem 8). The first part of this theorem is due to Bianchi [Bi1], the second part was proved in [GT] using Bianchi's analytic description of the Ribaucour transformations.
- In addition to orthogonal systems, we also treat, in a completely analogous manner, conjugate nets. We pose a Cauchy problem for them and show that any continuous conjugate net, along with its transformations, can be C^∞ -approximated by a sequence of discrete conjugate nets. The latter are quadrilateral lattices introduced by Doliwa and Santini [DS1].

The paper is organized as follows. We start in Sect. 2 with a short presentation of the concepts developed in our recent paper [BMS], stating the main result about an approximation of solutions of a hyperbolic system of partial differential equations by solutions of the corresponding partial difference system. In Sect. 3 we apply this theorem to conjugate systems. We formulate a Cauchy problem for smooth and discrete conjugate systems, define the Jonas transformation and prove the convergence. Sect. 4 is dedicated to the description of orthogonal systems in Möbius geometry. Sect. 5 contains the approximation results for orthogonal systems, a part of which was summarized above.

2 The approximation theorem

We start with a study of hyperbolic systems of partial difference and partial differential equations. A general convergence theorem is formulated, stating that solutions of difference equations approximate, under certain conditions, solutions of differential equations, provided the approximation of equations themselves takes place.

2.1 Notations

For a vector $\epsilon = (\epsilon_1, \dots, \epsilon_M)$ of M non-negative numbers, we define the lattice

$$\mathcal{P}^\epsilon = (\epsilon_1 \mathbb{Z}) \times \dots \times (\epsilon_M \mathbb{Z}) \subset \mathbb{R}^M. \quad (1)$$

We refer to ϵ_i as the *mesh size* of the grid $\mathcal{P}^\epsilon$ in the i -th direction; a direction with a vanishing mesh size $\epsilon_i = 0$ is understood as a *continuous* one, i.e. in this case the corresponding factor $\epsilon_i \mathbb{Z}$ in (1) is replaced by $\mathbb{R}$. For instance, the choice $\epsilon = (0, \dots, 0)$ yields $\mathcal{P}^\epsilon = \mathbb{R}^M$. We use also the following notation for n -dimensional sublattices of $\mathcal{P}^\epsilon$ ($n \leq M$):

$$\mathcal{P}_{i_1, \dots, i_n}^\epsilon = \{\xi = (\xi_1, \dots, \xi_M) \in \mathcal{P}^\epsilon \mid \xi_i = 0 \text{ if } i \notin \{i_1, \dots, i_n\}\}. \quad (2)$$

Considering convergence problems, we will mainly deal with the following two situations:

1. $\epsilon_i = \epsilon$ for all $i = 1, \dots, M$ (all directions become continuous in the limit $\epsilon \rightarrow 0$);
2. $\epsilon_i = \epsilon$ for $i = 1, \dots, m$; $\epsilon_i = 1$ for $i = m + 1, \dots, M$ (the last $M - m$ directions are kept discrete in the limit).

Thus, we always have one small parameter $\epsilon \geq 0$ only. In our geometric problems, the first situation corresponds to the approximation of conjugate and orthogonal systems themselves, while the second situation corresponds to the approximation of such systems along with their transformations. The underlying difference equations for discretizations of conjugate (resp. orthogonal) systems, and for their transformations will be the same, the only distinction being in the way the continuous limit is performed. We will distinguish between these two situations by setting $M = m + m'$, where $m' = 0$ and in the first situation, while $m' > 0$ in the second one.

Our considerations are local, therefore the encountered functions are usually defined on compact domains, like the cube

$$B(r) = \{\xi \in \mathbb{R}^M \mid 0 \leq \xi_i \leq r \text{ for } i \leq M\} \quad (3)$$

in the continuous context, and the following compact subsets of $\mathcal{P}^\epsilon$ in the lattice context. In the situation 1 ($m' = 0$):

$$B^\epsilon(r) = \{\xi \in \mathcal{P}^\epsilon \mid 0 \leq \xi_i \leq r \text{ for } i \leq M\}; \quad (4)$$

in the situation 2 ($m' > 0$):

$$B^\epsilon(r) = \{\xi \in \mathcal{P}^\epsilon \mid 0 \leq \xi_i \leq r \text{ for } 1 \leq i \leq m, \quad \xi_i \in \{0, 1\} \text{ for } m < i \leq M\}. \quad (5)$$

We will often use the notation

$$B_0^\epsilon(r) = \{\xi \in (\epsilon\mathbb{Z})^m \mid 0 \leq \xi_i \leq r \text{ for } 1 \leq i \leq m\}, \quad (6)$$

so that in the latter case $B^\epsilon(r) = B_0^\epsilon(r) \times \{0, 1\}^{m'}$.

For lattice functions $u : \mathcal{P}^\epsilon \mapsto \mathcal{X}$ with values in some Banach space $\mathcal{X}$, we use the shift operators τ_i and the partial difference operators $\delta_i = (\tau_i - \mathbf{1})/\epsilon_i$ defined as

$$(\tau_i u)(\xi) = u(\xi + \epsilon_i \mathbf{e}_i), \quad (\delta_i u)(\xi) = \frac{1}{\epsilon_i} (u(\xi + \epsilon_i \mathbf{e}_i) - u(\xi)); \quad (7)$$

here $\mathbf{e}_i$ is the i th unit vector. For $\epsilon_i = 0$, define $\tau_i = \mathbf{1}$ and $\delta_i = \partial_i$. Multiple shifts and higher order partial differences are written with the help of multi-indices $\alpha = (\alpha_1, \dots, \alpha_M) \in \mathbb{N}^M$:

$$\tau^\alpha = \tau_1^{\alpha_1} \cdots \tau_M^{\alpha_M}, \quad \delta^\alpha = \delta_1^{\alpha_1} \cdots \delta_M^{\alpha_M}, \quad |\alpha| = \alpha_1 + \dots + \alpha_M. \quad (8)$$

We use the following C^ℓ -norms for functions $u : B^\epsilon(r) \rightarrow \mathcal{X}$:

$$\|u\|_\ell = \sup \left\{ |\delta^\alpha u(\xi)|_{\mathcal{X}} \mid \xi \in B^\epsilon(r - \epsilon|\alpha|); |\alpha| \leq \ell \right\}; \quad (9)$$

if $m' > 0$, only the multi-indices α with $\alpha_i = 0$ ($i > m$) are taken into account in the last formula. Any function defined on $B^\epsilon(r)$ with $\epsilon > 0$ has finite C^ℓ -norms of arbitrary order ℓ ; on the other hand, if $u : B(r) \mapsto \mathcal{X}$ is a function of a continuous argument which does not belong to the class $C^\ell(B(r))$, then for its restriction to the lattice points, $u^\epsilon = u|_{B^\epsilon(r)}$, the norm $\|u^\epsilon\|_\ell$ will, in general, diverge for $\epsilon \rightarrow 0$.

Definition 1 *We say that a family of maps $\{u^\epsilon : B^\epsilon(r) \rightarrow \mathcal{X}\}_{0 < \epsilon < \epsilon_1}$ is $O(\epsilon)$ -convergent in $C^\infty(B(r))$ to a smooth limit $u : B(r) \rightarrow \mathcal{X}$, if there are finite real constants K_ℓ such that*

$$\|u^\epsilon - u|_{B^\epsilon(r)}\|_\ell \leq K_\ell \epsilon.$$

for any integer $\ell \geq 0$ and all positive $\epsilon < r/\ell$.

2.2 Approximation for hyperbolic equations

Being interested in the convergence of solutions of difference equations towards the solutions of differential ones, we restrict our attention to two versions of the limit behaviour of the lattice $\mathcal{P}^\epsilon$ described above, i.e. $m' = 0$ (all $\epsilon_i = \epsilon$) or $m' > 0$ ($\epsilon_i = \epsilon$ for $1 \leq i \leq m$, and $\epsilon_i = 1$ for $m+1 \leq i \leq M = m + m'$).

Let $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_K$ be a direct product of K Banach spaces, equipped with the norm

$$|u(\xi)|_{\mathcal{X}} = \max_{1 \leq k \leq K} |u_k(\xi)|_{\mathcal{X}_k}. \quad (10)$$

Consider a hyperbolic system of first-order partial difference ($\epsilon > 0$) or partial differential ($\epsilon = 0$) equations for functions $u : \mathcal{P}^\epsilon \mapsto \mathcal{X}$:

$$\delta_j u_k = f_{k,j}^\epsilon(u), \quad 1 \leq k \leq K, \quad j \in \mathcal{E}(k). \quad (11)$$

Thus, for each $k = 1, \dots, K$, there is defined a subset of indices $\mathcal{E}(k) \subset \{1, \dots, M\}$, called the set of *evolution directions* for the component u_k ; the complementary subset $\mathcal{S}(k) = \{1, \dots, M\} \setminus \mathcal{E}(k)$ is called the set of *static directions* for the component u_k . The dependent variables $u_k(\xi) = u_k(\xi_1, \dots, \xi_M)$ are thought of as attached to the cubes of dimension $\#(\mathcal{S}_k)$ adjacent to the point ξ .

The functions $f_{k,j}^\epsilon$ are defined on open domains $\mathbb{D}_{k,j}(\epsilon) \subset \mathcal{X}$ and are of class C^∞ there. We suppose that the following conditions are satisfied.

- (F) The domains $\mathbb{D}_{k,j}(\epsilon)$ for $1 \leq j \leq m$ (that is for all directions which become continuous in the limit $\epsilon \rightarrow 0$) exhaust $\mathcal{X}$ as $\epsilon \rightarrow 0$, i.e. any compact subset $\mathbb{K} \subset \mathcal{X}$ is contained in $\mathbb{D}_{k,j}(\epsilon)$ for all k , for all $j \in \mathcal{E}(k)$, $j \leq m$, and for all ϵ small enough. Similarly, there exists an open set $\mathbb{D} \subset \mathcal{X}$ such that the domains $\mathbb{D}_{k,j}(\epsilon)$ for $m+1 \leq j \leq M$ corresponding to the directions remaining discrete in the limit, exhaust $\mathbb{D}$ as $\epsilon \rightarrow 0$. The functions $f_{k,j}^\epsilon$ locally converge to the respective $f_{k,j}^0$ in $C^\ell(\mathcal{X})$, resp. in $C^\ell(\mathbb{D})$, for any ℓ , with the rate of convergence $O(\epsilon)$.

The existence of the subset $\mathbb{D}$ reflects the fact that transformations of conjugate, resp. orthogonal, systems are not defined for all initial data. The set $\mathbb{D}$ consists of “good” initial values, where the transformations are well defined.

Goursat Problem 1 *Given functions U_k^ϵ on $\mathcal{P}_{\mathcal{S}(k)}^\epsilon$ (the subspace spanned by the static directions of u_k), and $r > 0$, find a function $u^\epsilon : B^\epsilon(r) \rightarrow \mathcal{X}$ that solves the system (11) and satisfies the initial conditions $u_k^\epsilon = U_k^\epsilon$ on $\mathcal{P}_{\mathcal{S}(k)}^\epsilon \cap B^\epsilon(r)$.*

In order for this problem to admit a unique solution for $\epsilon > 0$, this has to be the case just for one elementary M -dimensional cube $B^\epsilon(\epsilon)$. The corresponding condition is called *consistency*, and it turns out to be necessary and sufficient also for the global (formal) solvability of the above Goursat problem. The consistency condition can be expressed as

$$\delta_j(\delta_i u_k) = \delta_i(\delta_j u_k), \quad (12)$$

for any choice of $i \neq j$ from the respective $\mathcal{E}_k$, if one writes out all the involved δu 's according to the equations (11). The first step in writing this out reads:

$$\delta_j(f_{k,i}(u)) = \delta_i(f_{k,j}(u)), \quad (13)$$

In order for the left-hand side (say) of this equation to be well-defined, the function $f_{k,i}$ is allowed to depend only on those u_ℓ for which $j \in \mathcal{E}_\ell$. This has to hold for all $j \in \mathcal{E}_k$, $j \neq i$, so we come to the condition that $f_{k,i}$ depends only on those u_ℓ for which $\mathcal{E}_k \setminus \{i\} \subset \mathcal{E}_\ell$. If this is satisfied, then the final form of the consistency condition reads:

$$\epsilon_j^{-1} \left(f_{k,i}(u + \epsilon_j f_j(u)) - f_{k,i}(u) \right) = \epsilon_i^{-1} \left(f_{k,j}(u + \epsilon_i f_i(u)) - f_{k,j}(u) \right), \quad (14)$$

where we use the abbreviation $f_i(u)$ for the K -vector whose k th component is $f_{k,i}(u)$, if $i \in \mathcal{E}_k$, and is not defined otherwise.

It is readily seen that, if one prescribes arbitrarily the values of all variables u_k at $\xi = 0$ (recall, $u_k(0)$ is actually associated with the respective static facet of dimension $\#(\mathcal{S}_k)$ adjacent to the point $\xi = 0$), then the system (11) defines uniquely the values for all variables on the remaining facets of the unit cube if and only if it is consistent. The following proposition shows that the local consistency assures also the global solvability of the Goursat problem; the only obstruction could appear if the solution would hit the set where the functions $f_{k,j}$ are not defined.

Proposition 1 *Let $\epsilon > 0$, and assume that the system (11) is consistent. Then either the Goursat problem 1 has a unique solution u^ϵ on the whole of $B^\epsilon(r)$, or there is a subdomain $\Omega \subset B^\epsilon(r)$ such that the Goursat problem 1 has a unique solution u^ϵ on Ω , and $u^\epsilon(\xi) \notin \mathbb{D}(\epsilon)$ at some point $\xi \in \Omega$.*

Finally, we turn to the whole family ($0 \leq \epsilon \leq \epsilon_0$) of Goursat problems for equations (11) with the respective data U_k^ϵ . The following result has been proven in our recent paper [BMS].

Theorem 2 *Let a family of Goursat problems 1 be given. Suppose that they are consistent for all $\epsilon > 0$, and satisfy the condition (F).*

1. *Let the functions $U_1^0, \dots, U_K^0$ be smooth, and, if $m' > 0$, assume that the point $(U_1^0(0), \dots, U_K^0(0))$ belongs to the subset $\mathbb{D}$. Then there exists $r > 0$ such that the Goursat problem 1 at $\epsilon = 0$ has a unique solution $u : B(r) \rightarrow \mathcal{X}$.*
2. *If the Goursat data U_k^ϵ locally $O(\epsilon)$ -converge in C^∞ to the functions U_k^0 , then there exists $\epsilon_1 \in (0, \epsilon_0)$ such that the Goursat problems 1 are solvable on $B^\epsilon(r)$ for all $0 < \epsilon < \epsilon_1$, and the solutions $u^\epsilon : B^\epsilon(r) \rightarrow \mathcal{X}$ converge to $u : B(r) \rightarrow \mathcal{X}$ with the rate $O(\epsilon)$ in $C^\infty(B(r))$.*

3 Conjugate nets

We give definitions of continuous and discrete conjugate nets and their transformations, formulate corresponding Goursat problems, and prove convergence of discrete conjugate nets towards continuous ones.

3.1 Basic definitions

Recall that $M = m + m'$. In Definition 2 below it is supposed that $m' = 0$, $M = m$, while in Definition 3 it is supposed that $m' = 1$, $M = m + 1$.

Definition 2 *A map $x : B_0(r) \mapsto \mathbb{R}^N$ is called an m -dimensional conjugate net in $\mathbb{R}^N$, if $\partial_i \partial_j x \in \text{span}(\partial_i x, \partial_j x)$ at any point $\xi \in B_0(r)$ for all pairs $1 \leq i \neq j \leq m$, i.e. if there exist functions $c_{ij}, c_{ji} : B_0(r) \mapsto \mathbb{R}$ such that*

$$\partial_i \partial_j x = c_{ji} \partial_i x + c_{ij} \partial_j x. \quad (15)$$

Definition 3 *A pair of m -dimensional conjugate nets $x, x^+ : B_0(r) \mapsto \mathbb{R}^N$ is called a Jonas pair, if three vectors $\partial_i x$, $\partial_i x^+$ and $\delta_M x = x^+ - x$ are coplanar at any point $\xi \in B_0(r)$ for all $1 \leq i \leq m$, i.e. if there exist functions $c_{iM}, c_{Mi} : B_0(r) \rightarrow \mathbb{R}$ such that*

$$\partial_i x^+ - \partial_i x = c_{Mi} \partial_i x + c_{iM} (x^+ - x). \quad (16)$$

Remarks.

- x^+ is also called a *Jonas transformation* of x . One can iterate these transformations and obtain sequences $(x^{(0)}, x^{(1)}, \dots, x^{(T)})$ of conjugate nets $x^{(t)} : B_0(r) \rightarrow \mathbb{R}^N$. It is natural to think of such sequences as of conjugate systems with m continuous and one discrete direction. We will see immediately that the notion of fully discrete conjugate systems puts all directions on an equal footing.
- A *Combescure transformation* x^+ of x is a Jonas transformation for which vectors $\partial_i x^+$ and $\partial_i x$ are parallel, $i = 1, \dots, m$. This class of transformations is singled out by requiring $c_{iM} = 0$ in equation (16).

Discrete conjugate nets were introduced by Doliwa and Santini [DS1].

Definition 4 *A map $x : B^\epsilon(r) \mapsto \mathbb{R}^N$ is called an M -dimensional discrete conjugate net in $\mathbb{R}^N$, if the four points x , $\tau_i x$, $\tau_j x$, and $\tau_i \tau_j x$ are coplanar at any $\xi \in B^\epsilon(r)$ for all pairs $1 \leq i \neq j \leq M$, i.e. if there exist functions $c_{ij} : B^\epsilon(r) \rightarrow \mathbb{R}$ such that*

$$\delta_i \delta_j x = c_{ji} \delta_i x + c_{ij} \delta_j x. \quad (17)$$

If $m' = 1$, $M = m + 1$, then the M -dimensional discrete conjugate net, considered as a pair of functions $x, x^+ : B_0^\epsilon(r) \mapsto \mathbb{R}^N$, is also called a Jonas pair of m -dimensional discrete conjugate nets.

3.2 Hyperbolic equations for conjugate nets

Introducing M new functions $w_i : B^\epsilon(r) \rightarrow \mathbb{R}^N$, we can rewrite (17) as system of first order:

$$\delta_i x = w_i, \quad (18)$$

$$\delta_i w_j = c_{ji} w_i + c_{ij} w_j, \quad i \neq j, \quad (19)$$

$$\delta_i c_{kj} = (\tau_j c_{ik}) c_{kj} + (\tau_j c_{ki}) c_{ij} - (\tau_i c_{kj}) c_{ij}, \quad i \neq j \neq k \neq i. \quad (20)$$

For a given discrete conjugate net $x : B^\epsilon(r) \rightarrow \mathbb{R}^N$, eq. (18) defines the functions w_j , then eq. (19) reflects the property (17), and eq. (20) is just a transcription of the compatibility condition $\delta_i(\delta_j w_k) = \delta_j(\delta_i w_k)$. Conversely, for any solution of (18)–(20) the map $x : B^\epsilon(r) \rightarrow \mathbb{R}^N$ has the defining property (17) and thus is a discrete conjugate net. So, discrete conjugate nets are in a one-to-one correspondence to solutions of the system (18)–(20).

This system almost suits the framework of Sect. 2.2; the only obstruction is the implicit nature of the equations (20) (their right-hand sides depend on the shifted variables like $\tau_i c_{kj}$ which is not allowed in (11). We return to this point later, and for a moment we handle the system (18)–(20) just as if it would belong to the class (11). In this context, we have to assign to every variable a Banach space and static/evolution directions. Abbreviating $w = (w_1, \dots, w_M)$ and $c = (c_{ij})_{i \neq j}$, we set

$$\mathcal{X} = \mathbb{R}^N \{x\} \times (\mathbb{R}^N)^M \{w\} \times \mathbb{R}^{M(M-1)} \{c\},$$

with the norm from (10). We assign to x no static directions, to w_i the only static direction i , and to c_{ij} two static directions i and j . The corresponding Goursat problem is now formulated as follows.

Goursat Problem 2 (for discrete conjugate nets). *Given:*

- a point $X^\epsilon \in \mathbb{R}^N$,
- M functions $W_i^\epsilon : \mathcal{P}_i^\epsilon \mapsto \mathbb{R}^N$ on the respective coordinate axes $\mathcal{P}_i^\epsilon$, and
- $M(M-1)$ functions $C_{ij}^\epsilon : \mathcal{P}_{ij}^\epsilon \mapsto \mathbb{R}$ on the coordinate planes $\mathcal{P}_{ij}^\epsilon$,

find a solution $(x^\epsilon, w^\epsilon, c^\epsilon) : B^\epsilon(r) \mapsto \mathcal{X}$ to the equations (18)–(20) satisfying the initial conditions

$$x^\epsilon(0) = X^\epsilon, \quad w_i^\epsilon|_{\mathcal{P}_i^\epsilon} = W_i^\epsilon, \quad c_{ij}^\epsilon|_{\mathcal{P}_{ij}^\epsilon} = C_{ij}^\epsilon. \quad (21)$$

We discuss now the consistency of the discrete hyperbolic system (18)–(20). First of all, note that for any triple of pairwise different indices (i, j, k) the equations of this system involving these indices only form a closed subset. In this sense, the system (18)–(20) consists of three-dimensional building blocks. One says also that it is *essentially three-dimensional*. Correspondingly, in the case $M = 3$ consistency is not an issue: all conditions to be verified are automatically taken into account by the very construction of the equations (18)–(20).

In this case $M = 3$ the initial data for an elementary cube are: a point $x(0)$, three vectors $w_1(0)$, $w_2(0)$, $w_3(0)$ which are thought of as attached to the edges incident to the point $\xi = 0$ and parallel to the correspondent axes, and six numbers $c_{ij}(0)$, $1 \leq i \neq j \leq 3$, which are thought of as attached to the plaquettes incident to the point $\xi = 0$ and parallel to the coordinate planes $\mathcal{P}_{ij}$. From these data, one constructs first with the help of (18) the three points $\tau_i x = x(0) + \epsilon_i w_i(0)$, then one calculates $\tau_i w_j$ for $i \neq j$ with the help of (19), and further $\tau_j \tau_i x$ by means of (18). Notice that the consistency $\tau_j \tau_i x = \tau_i \tau_j x$ is assured by the symmetry of the right-hand side of (19) with respect to the indices i and j . Thus, the first two equations (18), (19) allow us to determine the values of x in seven vertices of an elementary cube. The remaining 8-th one, $\tau_1 \tau_2 \tau_3 x$, is now determined as follows. Calculate $\tau_i c_{kj}$ from the (linearly implicit) equations (20), then calculate $\tau_j \tau_i w_k$ from (19), and then $\tau_1 \tau_2 \tau_3 x$ from (18). Again, the consistency requirement $\tau_j \tau_i w_k = \tau_i \tau_j w_k$ is guaranteed, per construction, by eqs. (20). Geometrically, the need to solve the linearly implicit equations for $\tau_i c_{kj}$ is interpreted as follows. The point $\tau_1 \tau_2 \tau_3 x$ is determined by the conditions that it lies in three planes $\tau_i \Pi_{jk}$, $1 \leq i \leq 3$, where $\tau_i \Pi_{jk}$ is defined as the plane passing through three points $(\tau_i x, \tau_j \tau_i x, \tau_k \tau_i x)$, or, equivalently, passing through $\tau_i x$ and spanned by $\tau_i w_j$, $\tau_i w_k$. All three planes $\tau_1 \Pi_{23}$, $\tau_2 \Pi_{13}$ and $\tau_3 \Pi_{12}$ belong to the three-dimensional space through $x(0)$ spanned by $w_i(0)$, $1 \leq i \leq 3$, and therefore generically they intersect at exactly one point. The above mentioned linearly implicit system encodes just finding the intersection point of three planes.

Turning now to the case $M > 3$, we face a non-trivial consistency problem. Indeed, the point $\tau_i \tau_j \tau_k \tau_\ell x$ can be constructed as the intersection point of three planes $\tau_i \tau_j \Pi_{k\ell}$, $\tau_i \tau_k \Pi_{j\ell}$, $\tau_i \tau_\ell \Pi_{jk}$, where the plane $\tau_i \tau_j \Pi_{k\ell}$ is defined as the plane passing through three points $(\tau_i \tau_j x, \tau_i \tau_j \tau_k x, \tau_i \tau_j \tau_\ell x)$, or, equivalently, passing through $\tau_i \tau_j x$ and spanned by $\tau_i \tau_j w_k$, $\tau_i \tau_j w_\ell$. So, there are four alternative ways to determine the point $\tau_i \tau_j \tau_k \tau_\ell x$, depending on which index plays the role of i .

Proposition 2 *The system (18)–(20) is consistent for $M > 3$.*

Proof. The geometric meaning of this statement is that the above mentioned four ways to obtain $\tau_i\tau_j\tau_k\tau_\ell x$ lead to one and the same result. This can be proved by symbolic manipulations with the equations (18)–(20), but a geometric proof first given in [DS1] is much more transparent. In the construction above, it is easy to understand that the plane $\tau_i\tau_j\Pi_{k\ell}$ is the intersection of two three-dimensional subspaces $\tau_i\Pi_{jkl}$ and $\tau_j\Pi_{ikl}$ in the four-dimensional space through $x(0)$ spanned by $w_i(0)$, $w_j(0)$, $w_k(0)$, $w_\ell(0)$, where the subspace $\tau_i\Pi_{jkl}$ is the one through the four points $(\tau_i x, \tau_i\tau_j x, \tau_i\tau_k x, \tau_i\tau_\ell x)$, or, equivalently, the one through $\tau_i x$ spanned by $\tau_i w_j$, $\tau_i w_k$ and $\tau_i w_\ell$. Now the point $\tau_i\tau_j\tau_k\tau_\ell x$ can be alternatively described as the unique intersection point of the four three-dimensional subspaces $\tau_i\Pi_{jkl}$, $\tau_j\Pi_{ikl}$, $\tau_k\Pi_{ijl}$ and $\tau_\ell\Pi_{ijk}$ of one and the same four-dimensional space. ■

Now, we turn to the above mentioned feature of the system (18)–(20), namely to its implicit nature. The geometric background of this feature is the following. Though there is, in general, a unique way to construct the 8-th vertex $\tau_i\tau_j\tau_k x$ of an elementary hexahedron out of the the seven vertices x , $\tau_i x$ and $\tau_i\tau_j x$, or, equivalently, out of x , w_i , c_{ij} , it can happen that the resulting hexahedron is degenerate. This happens, e.g., if the point $\tau_i\tau_j x$ lies in the plane $\tau_k\Pi_{ij}$ through $(\tau_k x, \tau_i\tau_k x, \tau_j\tau_k x)$. This is illustrated on Fig. 4. While the example on the left of Fig. 4 is fine, the right one does not



Figure 4: A non-degenerate and a degenerate hexahedron.

have the combinatorics of a 3-cube (the top-right edge has degenerated to a point). As a result, some of the quantities $\tau_i c_{kj}$ in equation (20) are not well-defined; this is reflected by the fact that the equations (20) are implicit and have to be solved for $\tau_i c_{kj}$, which is possible not for all values of c but rather for those belonging to a certain open subset.

To investigate this point, rewrite (20) as

$$\delta_i c_{kj} = F_{(ijk)}(c) + \epsilon_j(\delta_j c_{ik})c_{kj} + \epsilon_j(\delta_j c_{ki})c_{ij} - \epsilon_i(\delta_i c_{kj})c_{ij} \quad (22)$$

Introducing vectors δc and $F(c)$ with $M(M-1)(M-2)$ components labelled

by triples (ijk) of pairwise distinct numbers $1 \leq i, j, k \leq M$,

$$\begin{aligned}\delta c_{(ijk)} &= \delta_i c_{kj} \\ F_{(ijk)}(c) &= c_{ik}c_{kj} + c_{ki}c_{ij} - c_{kj}c_{ij},\end{aligned}$$

we present the above equations as

$$(\mathbf{1} - Q^\epsilon(c))\delta c = F(c), \quad (23)$$

with a suitable matrix $Q^\epsilon(c)$. We need to find conditions for $\mathbf{1} - Q^\epsilon(c)$ to be invertible.

3.2.1 One conjugate net

If $m' = 0$, then all entries of $Q^\epsilon(c)$ are either zero or of the form ϵc_{ij} . Therefore, on any compact $\mathbb{K} \subset \mathcal{X}$ we have: $\|Q^\epsilon(c)\| < C\epsilon$ with some $C = C(\mathbb{K}) > 0$, hence $\mathbf{1} - Q^\epsilon(c)$ is invertible for $\epsilon < \epsilon_0 = 1/C$, and the inverse matrix $(\mathbf{1} - Q^\epsilon(c))^{-1}$ is a smooth function on $\mathbb{K}$ and $O(\epsilon)$ -convergent in $C^\infty(\mathbb{K})$ to $\mathbf{1}$. So, the system (23) is solvable, and

$$\delta_i c_{kj} = F_{(ijk)}(c) + O(\epsilon) = c_{ik}c_{kj} + c_{ki}c_{ij} - c_{kj}c_{ij} + O(\epsilon),$$

where the constant in $O(\epsilon)$ is uniform on any compact set.

The limiting equations for (18)-(20) in this case are:

$$\partial_i x = w_i \quad (24)$$

$$\partial_i w_j = c_{ji}w_i + c_{ij}w_j \quad (25)$$

$$\partial_i c_{kj} = c_{ik}c_{kj} + c_{ki}c_{ij} - c_{kj}c_{ij}. \quad (26)$$

For any solution $(x, w, c) : B_0(r) \rightarrow \mathcal{X}$ of this system the function $x : B_0(r) \rightarrow \mathbb{R}^N$ is a conjugate net, because equations (24) and (25) yield the defining property (15). Conversely, if $x : B_0(r) \rightarrow \mathbb{R}^N$ is a conjugate net, then equation (24) defines the vectors w_i , these fulfill (25) since this is the defining property of a conjugate net, and (26) results from equating $\partial_i(\partial_j w_k) = \partial_j(\partial_i w_k)$. So, the following is demonstrated:

Lemma 1 *If $m' = 0$, then the Goursat problem 2 for discrete conjugate nets fulfills the condition (F). The limiting system for $\epsilon = 0$, consisting of (24)-(26), describes continuous conjugate nets.*

3.2.2 Jonas pair of conjugate nets

In the case $m' = 1$ the matrix $\mathbf{1} - Q^\epsilon(c)$ is no longer a small perturbation of the identity, since $\epsilon_M = 1$. More precisely, the matrix $Q^\epsilon(c)$ is block-diagonal, its $\binom{M}{3}$ diagonal blocks $Q_{\{ijk\}}^\epsilon(c)$ of size 6×6 correspond to non-ordered triples of pairwise distinct indices $\{i, j, k\}$, with rows and columns labeled by the six possible permutations of these indices. Blocks for which $i, j, k \neq M$ are of the form considered before – their entries are either zero or of the form ϵc_{ij} . Blocks where, say, $k = M$, admit a decomposition $Q_{\{ijM\}}^\epsilon(c) = A(c) + \epsilon B(c)$, where the matrices A and B do not depend on ϵ . It is not difficult to calculate that $\det(\mathbf{1} - A(c)) = (1 + c_{Mi})(1 + c_{Mj})$. So $\mathbf{1} - Q_{\{ijM\}}^\epsilon(c)$ is invertible if $c_{Mi}, c_{Mj} \neq -1$, provided ϵ is small enough. The inverse is smooth in a neighborhood any such point c , and $O(\epsilon)$ -convergent to $(\mathbf{1} - A(c))^{-1}$.

As formal limit of (18) and (19), we obtain ($i \neq j$):

$$\partial_i x = w_i \quad (1 \leq i \leq m), \quad (27)$$

$$\delta_M x = w_M, \quad (28)$$

$$\partial_i w_j = c_{ji} w_i + c_{ij} w_j \quad (1 \leq i \leq m), \quad (29)$$

$$\delta_M w_j = c_{jM} w_M + c_{Mj} w_j. \quad (30)$$

There are also four different limits of equation (20), depending on which directions i and j are kept discrete. We shall not write them down; they are easily reconstructed as the compatibility conditions for (29), (30), like $\delta_M(\partial_i w_k) = \partial_i(\delta_M w_k)$ etc. Comparing (27)–(30) with the definition of the Jonas transformation (16), we see that the following is demonstrated.

Lemma 2 *If $m' = 1$, then the Goursat problem 2 for discrete conjugate nets also fulfills the condition (F), with the domain*

$$\mathbb{D} = \{(x, w, c) \in \mathcal{X} \mid c_{Mi} \neq -1 \text{ for } 1 \leq i \leq m\}.$$

The limiting system at $\epsilon = 0$ describes Jonas pairs of continuous conjugate nets.

3.3 Approximation theorems for conjugate nets

We are now ready to formulate and prove the main results of this chapter.

Theorem 3 (approximation of a conjugate net). *Let there be given:*

- m smooth curves $X_i : \mathcal{P}_i \mapsto \mathbb{R}^N$, $i = 1, \dots, m$, intersecting at a common point $X = X_1(0) = \dots = X_m(0)$;

- for each pair $1 \leq i < j \leq m$, two smooth functions $C_{ij}, C_{ji} : \mathcal{P}_{ij} \mapsto \mathbb{R}$.

Then, for some $r > 0$:

1. There is a unique conjugate net $x : B_0(r) \rightarrow \mathbb{R}^N$ such that

$$x|_{\mathcal{P}_i} = X_i, \quad c_{ij}|_{\mathcal{P}_{ij}} = C_{ij}. \quad (31)$$

2. The family of discrete conjugate nets $\{x^\epsilon : B_0^\epsilon(r) \rightarrow \mathbb{R}^N\}_{0 < \epsilon < \epsilon_1}$ uniquely determined by requiring

$$x^\epsilon|_{\mathcal{P}_i^\epsilon} = X_i|_{\mathcal{P}_i^\epsilon}, \quad c_{ij}^\epsilon|_{\mathcal{P}_{ij}^\epsilon} = C_{ij}|_{\mathcal{P}_{ij}^\epsilon}, \quad (32)$$

where c_{ij}^ϵ are the rotation coefficients of the net x^ϵ , $O(\epsilon)$ -converges in $C^\infty(B_0(r))$ to x .

Proof. It is easy to reformulate the data (31), (32) into valid initial data for the Goursat problem on $\mathcal{X}$ for the systems (24)–(26), resp. (18)–(20). Namely, at $\epsilon = 0$ one translates (31) into the equivalent data

$$x(0) = X, \quad w_i|_{\mathcal{P}_i} = \partial_i X_i, \quad c_{ij}|_{\mathcal{P}_{ij}} = C_{ij}. \quad (33)$$

while for $\epsilon > 0$ one translates (32) into the equivalent data

$$x^\epsilon(0) = X, \quad w_i^\epsilon|_{\mathcal{P}_i^\epsilon} = \delta_i X_i|_{\mathcal{P}_i^\epsilon}, \quad c_{ij}^\epsilon|_{\mathcal{P}_{ij}^\epsilon} = C_{ij}|_{\mathcal{P}_{ij}^\epsilon}. \quad (34)$$

Now the theorem follows directly from Lemma 1 and Theorem 2. ■

Theorem 4 (approximation of a Jonas pair). *Let, in addition to the data listed in Theorem 3, there be given:*

- m smooth curves $X_i^+ : \mathcal{P}_i \rightarrow \mathbb{R}^N$, $i = 1, \dots, m$, such that all X_i^+ intersect at a common point $X^+ = X_1^+(0) = \dots = X_m^+(0)$, and such that for any $1 \leq i \leq m$ and for any point $\xi \in \mathcal{P}_i$ the three vectors $\partial_i X_i$, $\partial_i X_i^+$ and $\delta_M X_i = X_i^+ - X_i$ are coplanar.

Assume that $X^+ - X$ is not parallel to any of the m vectors $\partial_i X_i^+(0)$. Define the functions $C_{Mi}, C_{iM} : \mathcal{P}_i \mapsto \mathbb{R}$ by the formula

$$\partial_i X_i^+ - \partial_i X_i = C_{Mi} \partial_i X_i + C_{iM} (X_i^+ - X_i). \quad (35)$$

Then, for some $r > 0$:

1. In addition to the conjugate net $x : B_0(r) \rightarrow \mathbb{R}^N$ defined in Theorem 3, there is its unique Jonas transformation $x^+ : B_0(r) \rightarrow \mathbb{R}^N$ such that

$$x^+|_{\mathcal{P}_i} = X_i^+ \quad (1 \leq i \leq m); \quad (36)$$

2. The family $\{x^\epsilon : B^\epsilon(r) \rightarrow \mathbb{R}^N\}_{0 < \epsilon < \epsilon_1}$ of discrete M -dimensional conjugate nets ($M = m + 1$), uniquely determined by the requirements

$$x^\epsilon|_{\mathcal{P}_i^\epsilon} = X_i|_{\mathcal{P}_i^\epsilon}, \quad c_{ij}^\epsilon|_{\mathcal{P}_{ij}^\epsilon} = C_{ij}|_{\mathcal{P}_{ij}^\epsilon}, \quad (1 \leq i \neq j \leq m) \quad (37)$$

and

$$x^\epsilon(\mathbf{0}, 1) = X^+, \quad c_{Mi}^\epsilon|_{\mathcal{P}_i^\epsilon} = C_{Mi}|_{\mathcal{P}_i^\epsilon}, \quad c_{iM}^\epsilon|_{\mathcal{P}_i^\epsilon} = C_{iM}|_{\mathcal{P}_i^\epsilon} \quad (1 \leq i \leq m) \quad (38)$$

$O(\epsilon)$ -converges in $C^\infty(B(r))$ to the Jonas pair x, x^+ , in the sense that $x^\epsilon(\cdot, 0) \rightarrow x$, $x^\epsilon(\cdot, 1) \rightarrow x^+$

Proof. Consider the Goursat problem for M -dimensional discrete conjugate nets with the initial data

$$x^\epsilon(0) = X, \quad w_i^\epsilon|_{\mathcal{P}_i^\epsilon} = \delta_i X_i|_{\mathcal{P}_i^\epsilon}, \quad w_M^\epsilon(0) = X^+ - X, \quad (i = 1, \dots, m) \quad (39)$$

and

$$c_{ij}^\epsilon|_{\mathcal{P}_{ij}^\epsilon} = C_{ij}|_{\mathcal{P}_{ij}^\epsilon}, \quad c_{Mi}^\epsilon|_{\mathcal{P}_i^\epsilon} = C_{Mi}|_{\mathcal{P}_i^\epsilon}, \quad c_{iM}^\epsilon|_{\mathcal{P}_i^\epsilon} = C_{iM}|_{\mathcal{P}_i^\epsilon} \quad (1 \leq i \neq j \leq m). \quad (40)$$

The statement of the theorem follows by applying Theorem 2 to this situation. This, in turn, is possible due to Lemma 2 and the following observation. The condition of the theorem yields that $X_i^+(\xi) - X_i(\xi)$ is not parallel to any of $\partial_i X_i^+(\xi)$ not only at $\xi = 0$, but also in some neighborhoods of zero on the corresponding axes $\mathcal{P}_i$. Now one deduces from the definition (35) of the rotation coefficients C_{iM} , C_{Mi} that in the same neighborhoods $C_{Mi} \neq -1$. Hence, the data of our Goursat problem belong to the set $\mathbb{D}$ of Lemma 2. ■

Letting not one but two or three directions of the conjugate net remain discrete in the continuous limit (so that $m' = 2$ or $m' = 3$), one arrives at the permutability properties of the Jonas transformations.

Theorem 5 (permutability of Jonas transformations).

1. Given an m -dimensional conjugate net $x(\cdot, 0, 0) : B_0(r) \mapsto \mathbb{R}^N$ and its two Jonas transformations $x(\cdot, 1, 0) : B_0(r) \mapsto \mathbb{R}^N$ and $x(\cdot, 0, 1) : B_0(r) \mapsto \mathbb{R}^N$, there exists a two-parameter family of conjugate nets $x(\cdot, 1, 1) : B_0(r) \mapsto \mathbb{R}^N$ that are Jonas transformations of both $x(\cdot, 1, 0)$ and $x(\cdot, 0, 1)$. Corresponding points of the four conjugate nets are coplanar.

2. *Given three Jonas transformations*

$$x(\cdot, 1, 0, 0), x(\cdot, 0, 1, 0), x(\cdot, 0, 0, 1) : B_0(r) \mapsto \mathbb{R}^N$$

of a given m -dimensional conjugate net $x(\cdot, 0, 0, 0) : B_0(r) \mapsto \mathbb{R}^N$, as well as three further conjugate nets

$$x(\cdot, 1, 1, 0), x(\cdot, 0, 1, 1), x(\cdot, 1, 0, 1) : B_0(r) \mapsto \mathbb{R}^N$$

such that $x(\cdot, 1, 1, 0)$ is a Jonas transformation of both $x(\cdot, 1, 0, 0)$ and $x(\cdot, 0, 1, 0)$ etc., there exists generically a unique conjugate net

$$x(\cdot, 1, 1, 1) : B_0(r) \mapsto \mathbb{R}^N$$

which is a Jonas transformation of all three $x(\cdot, 1, 1, 0)$, $x(\cdot, 0, 1, 1)$ and $x(\cdot, 1, 0, 1)$.

These results are proved in the exactly same manner as Theorem 4, if one takes into account the multi-dimensional consistency of discrete conjugate nets (Proposition 2).

4 Orthogonal systems

We give the definitions of orthogonal systems and their transformations, formulate corresponding Goursat problems, and prove convergence of discrete orthogonal systems to continuous ones.

4.1 Basic definitions

Definition 5 *A conjugate net $x : B_0(r) \mapsto \mathbb{R}^N$ is called an m -dimensional orthogonal system in $\mathbb{R}^N$ (an orthogonal coordinate system if $m = N$), if*

$$\partial_i x \cdot \partial_j x = 0. \quad (41)$$

Definition 6 *A pair of m -dimensional orthogonal systems $x, x^+ : B_0(r) \mapsto \mathbb{R}^N$ is called a Ribaucour pair, if for any $1 \leq i \leq m$ the corresponding coordinate lines of x and x^+ envelope a one-parameter family of circles, i.e. if it is a Jonas pair of conjugate nets and*

$$\left(\frac{\partial_i x^+}{|\partial_i x^+|} + \frac{\partial_i x}{|\partial_i x|} \right) \cdot (x^+ - x) = 0. \quad (42)$$

Remarks.

- x^+ is also called a Ribaucour transformation of x ; Ribaucour transformations can be iterated, and our results generalize to any finite sequence of such transformations (cf. the remarks after Definition 3).
- Two-dimensional ($m = 2$) orthogonal systems are called *C-surfaces*. C-surfaces in $\mathbb{R}^3$ are characterized as surfaces parametrized along curvature lines.
- There are different definitions of Ribaucour transformations in the literature (cf. [H]). We have chosen the one that is best suited for our present purposes.

Definition 7 *A map $x: B^\epsilon(r) \mapsto \mathbb{R}^N$ is called an M -dimensional discrete orthogonal system in $\mathbb{R}^N$, if the four points $x, \tau_i x, \tau_j x$ and $\tau_i \tau_j x$ are concircular for all $\xi \in B^\epsilon(r)$ and all $1 \leq j \neq i \leq M$.*

If $m' = 1$, $M = m + 1$, then the M -dimensional discrete orthogonal system, considered as a pair of functions $x, x^+ : B_0^\epsilon(r) \mapsto \mathbb{R}^N$, is called a Ribaucour pair of m -dimensional discrete orthogonal systems.

Remark.

- Also in the discrete case we shall use the term *C-surfaces* for two-dimensional ($m = 2$) orthogonal systems.

So, orthogonal systems form a subclass of conjugate nets subject to a certain additional condition (orthogonality of coordinate lines, resp. circularity of elementary quadrilaterals).

The classical description of continuous orthogonal systems is this (in the following equations it is assumed that $i \neq j \neq k \neq i$):

$$\partial_i x = h_i v_i, \quad (43)$$

$$\partial_i v_j = \beta_{ji} v_i, \quad (44)$$

$$\partial_i h_j = h_i \beta_{ij}, \quad (45)$$

$$\partial_i \beta_{kj} = \beta_{ki} \beta_{ij}, \quad (46)$$

$$\partial_i \beta_{ij} + \partial_j \beta_{ji} = -\partial_i v_i \cdot \partial_j v_j. \quad (47)$$

Eq. (43) defines a system $\{v_i\}_{1 \leq i \leq m}$ of m orthonormal vectors at each point $\xi \in B_0(r)$; the quantities $h_i = |\partial_i x|$ are the respective metric coefficients. Conjugacy of the net x and normalization of v_i imply that eq. (44) holds with some real-valued functions β_{ij} . Eq. (45) expresses the consistency condition $\partial_i(\partial_j x) = \partial_j(\partial_i x)$ for $i \neq j$. Analogously, eq. (46) expresses the consistency condition $\partial_i(\partial_j v_k) = \partial_j(\partial_i v_k)$ for $i \neq j \neq k \neq i$. The m equations

(45) and $\binom{m}{3}$ equations (46) are called the *Darboux system*; they constitute a description of conjugate nets alternative to the system (24)–(26) we used in Sect. 3. The orthogonality constraint is expressed by $\binom{m}{2}$ additional equations (47), derived from the identity $\partial_i \partial_j \langle v_i, v_j \rangle = 0$. In the case $m = N$ the scalar product on the right-hand side of (47) can be expressed in terms of rotation coefficients β_{ij} only:

$$\partial_i \beta_{ij} + \partial_j \beta_{ji} + \sum_{k \neq i, j} \beta_{ki} \beta_{kj} = 0. \quad (48)$$

The Darboux system (45), (46) together with the eqs. (48) forms the *Lamé system*.

The classical approach as presented is based on the Euclidean geometry. However, the invariance group of orthogonal systems is the Möbius group, which acts on the compactification $\mathbb{R}^N \cup \{\infty\} \approx \mathbb{S}^N$, rather than on $\mathbb{R}^N$. This is a motivation to consider orthogonal systems in the sphere $\mathbb{S}^N$ and to study their Möbius-invariant description. In particular, this enables one to give a frame description of orthogonal systems which can be generalized to the discrete context in a straightforward manner. This turns out to be the key to derivation of the discrete analogue of the Lamé system.

4.2 Möbius geometry

We give a brief presentation of Möbius geometry, tailored for our current needs. A more profound introduction may be found in [H].

The N -dimensional Möbius geometry is associated with the unit sphere $\mathbb{S}^N \subset \mathbb{R}^{N+1}$. Fix two antipodal points on $\mathbb{S}^N$, $p_0 = (0, \dots, 0, 1)$ and $p_\infty = -p_0$. The standard stereographic projection σ from the point p_∞ is a conformal bijection from $\mathbb{S}_*^N = \mathbb{S}^N \setminus \{p_\infty\}$ to $\mathbb{R}^N$ that maps spheres (of any dimension) in $\mathbb{S}^N$ to spheres or affine subspaces in $\mathbb{R}^N$. Its inverse map

$$\sigma^{-1}(x) = \frac{2}{1 + |x|^2} x + \frac{1 - |x|^2}{1 + |x|^2} p_0$$

lifts an orthogonal system in $\mathbb{R}^N$, continuous or discrete, to an orthogonal system in $\mathbb{S}^N$.

The group $\mathcal{M}(N)$ of N -dimensional Möbius transformations consists of those bijective maps $\mu : \mathbb{S}^N \rightarrow \mathbb{S}^N$ that map any sphere in $\mathbb{S}^N$ to a sphere of the same dimension; it then follows that μ is also conformal.

Fact 1 *The notion of orthogonal system on $\mathbb{S}^N$ is invariant under Möbius transformations.*

Möbius transformations can be embedded into a matrix group, namely the group of pseudo-orthogonal transformations on the Minkowski space $\mathbb{R}^{N+1,1}$, i.e. the $(N+2)$ -dimensional space spanned by $\{\mathbf{e}_1, \dots, \mathbf{e}_{N+2}\}$ and equipped with the Lorentz scalar product

$$\left\langle \sum_{i=1}^{N+2} u_i \mathbf{e}_i, \sum_{j=1}^{N+2} v_j \mathbf{e}_j \right\rangle = \sum_{k=1}^{N+1} u_k v_k - u_{N+2} v_{N+2}.$$

To do this, a model is used where points of $\mathbb{S}^N$ are identified with lines on a *light cone*

$$\mathcal{L}^{N+1} = \{u \in \mathbb{R}^{N+1,1} \mid \langle u, u \rangle = 0\} \subset \mathbb{R}^{N+1,1}.$$

This identification is achieved via the projection map

$$\begin{aligned} \pi : \mathbb{R}^{N+1,1} \setminus (\mathbb{R}^{N+1} \times \{0\}) &\mapsto \mathbb{R}^{N+1}, \\ (u_1, \dots, u_{N+1}, u_{N+2}) &\mapsto (u_1/u_{N+2}, \dots, u_{N+1}/u_{N+2}), \end{aligned}$$

which sends lines on $\mathcal{L}^{N+1}$ to points of $\mathbb{S}^N$. In particular, the lines through

$$\mathbf{e}_0 = \frac{1}{2}(\mathbf{e}_{N+2} + \mathbf{e}_{N+1}) \quad \text{and} \quad \mathbf{e}_\infty = \frac{1}{2}(\mathbf{e}_{N+2} - \mathbf{e}_{N+1})$$

are mapped to the points p_0 and p_∞ , respectively.

The group $O^+(N+1, 1)$ of genuine Lorentz transformations consists of pseudo-orthogonal linear maps L which preserve “the direction of time”:

$$\langle L(u), L(v) \rangle = \langle u, v \rangle \quad \forall u, v \in \mathbb{R}^{N+1,1}, \quad \langle L(\mathbf{e}_{N+2}), \mathbf{e}_{N+2} \rangle < 0. \quad (49)$$

For $L \in O^+(N+1, 1)$, the projection π induces a Möbius transformation of $\mathbb{S}^N$ via $\mu(L) = \pi \circ L \circ \pi^{-1}$. Indeed, L is linear and preserves the light cone because of (49), so $\mu(L)$ maps $\mathbb{S}^N$ to itself. Further, L maps linear planes to linear planes; under π , linear planes in $\mathbb{R}^{N+1,1}$ project to affine planes in $\mathbb{R}^{N+1}$. Since any sphere in $\mathbb{S}^N$ is uniquely represented as the intersection of an affine plane in $\mathbb{R}^{N+1}$ with $\mathbb{S}^N \subset \mathbb{R}^{N+1}$, we conclude that spheres of any dimension are mapped by $\mu(L)$ to spheres of the same dimension, which is the defining property of Möbius transformations. It can be shown that the correspondence $L \mapsto \mu(L)$ between $L \in O^+(N+1, 1)$ and $\mathcal{M}(N)$ is indeed one-to-one.

Recall that we are interested in orthogonal systems in $\mathbb{R}^N$; their lifts to $\mathbb{S}^N$ actually do not contain p_∞ . Therefore, we focus on those Möbius transformations which preserve p_∞ , thus corresponding to Euclidean motions and homotheties in $\mathbb{R}^N = \sigma(\mathbb{S}_*^N)$. For these, we can restrict our attention to the section

$$\mathcal{K} = \left\{ u \in \mathbb{R}^{N+1,1} : \langle u, u \rangle = 0, \langle u, \mathbf{e}_\infty \rangle = -1/2 \right\} = \mathcal{L}^{N+1} \cap (\mathbf{e}_0 + \mathbf{e}_\infty^\perp).$$

(We use here the notation $u^\perp$ for the hyperplane orthogonal to u .) The canonical lift

$$\begin{aligned}\lambda : \mathbb{R}^N &\mapsto \mathcal{K}, \\ x &\mapsto x + \mathbf{e}_0 + |x|^2 \mathbf{e}_\infty,\end{aligned}$$

factors the (inverse) stereographic projection, $\sigma^{-1} = \pi \circ \lambda$, and is an isometry in the following sense: for any four points $x_1, \dots, x_4 \in \mathbb{R}^N$ one has:

$$\langle \lambda(x_1) - \lambda(x_2), \lambda(x_3) - \lambda(x_4) \rangle = (x_1 - x_2) \cdot (x_3 - x_4). \quad (50)$$

We summarize these results in a diagram

$$\begin{array}{ccc}\mathcal{K} & \hookrightarrow & \mathcal{L}^{N+1} \subset \mathbb{R}^{N+1,1} \\ \lambda \uparrow & & \downarrow \pi \\ \mathbb{R}^N & \xrightarrow{\sigma^{-1}} & \mathbb{S}^N \subset \mathbb{R}^{N+1}\end{array}$$

Denote by $O_\infty^+(N+1, 1)$ the subgroup of Lorentz transformations fixing the vector $\mathbf{e}_\infty$ (and consequently, preserving $\mathcal{K}$). The same arguments as before allow one to identify this subgroup with the group $E(N)$ of Euclidean motions of $\mathbb{R}^N$.

Fact 2 *Möbius transformations of $\mathbb{S}^N$ are in a one-to-one correspondence with the genuine Lorentz transformations of $\mathbb{R}^{N+1,1}$. Euclidean motions of $\mathbb{R}^N$ are in a one-to-one correspondence with the Lorentz transformations that fix $\mathbf{e}_\infty$.*

Our next technical device will be Clifford algebras which are very convenient to describe Lorentz transformations in $\mathbb{R}^{N+1,1}$, and hence Möbius transformations in $\mathbb{S}^N$. Recall that the Clifford algebra $\mathcal{C}(N+1, 1)$ is an algebra over $\mathbb{R}$ with generators $\mathbf{e}_1, \dots, \mathbf{e}_{N+2} \in \mathbb{R}^{N+1,1}$ subject to the relations

$$uv + vu = -2\langle u, v \rangle \mathbf{1} = -2\langle u, v \rangle \quad \forall u, v \in \mathbb{R}^{N+1,1}. \quad (51)$$

Relation (51) implies that $u^2 = -\langle u, u \rangle$, so any vector $u \in \mathbb{R}^{N+1,1} \setminus \mathcal{L}^{N+1}$ has an inverse $u^{-1} = -u/\langle u, u \rangle$. The multiplicative group generated by the invertible vectors is called the *Clifford group*. It contains the subgroup $Pin(N+1, 1)$ which is a universal cover of the group of Möbius transformations. We shall need the *genuine Pin group*:

$$\mathcal{H} = Pin^+(N+1, 1) = \{u_1 \cdots u_n \mid u_i^2 = -1\},$$

and its subgroup generated by vectors orthogonal to $\mathbf{e}_\infty$:

$$\mathcal{H}_\infty = \{u_1 \cdots u_n \mid u_i^2 = -1, \langle u_i, \mathbf{e}_\infty \rangle = 0\} \subset \mathcal{H}.$$

$\mathcal{H}$ and $\mathcal{H}_\infty$ are Lie groups with Lie algebras

$$\mathfrak{h} = \text{spin}(N+1, 1) = \text{span}\left\{\mathbf{e}_i\mathbf{e}_j : i, j \in \{0, 1, \dots, N, \infty\}, i \neq j\right\}, \quad (52)$$

$$\mathfrak{h}_\infty = \text{spin}_\infty(N+1, 1) = \text{span}\left\{\mathbf{e}_i\mathbf{e}_j : i, j \in \{1, \dots, N, \infty\}, i \neq j\right\}. \quad (53)$$

In fact, $\mathcal{H}$ and $\mathcal{H}_\infty$ are universal covers of the previously defined Lorentz subgroups $O^+(N+1, 1)$ and $O_\infty^+(N+1, 1)$, respectively. The covers are double, since the lift $\psi(L)$ of a Lorentz transformation L is defined up to a sign. To show this, consider the coadjoint action of $\psi \in \mathcal{H}$ on $v \in \mathbb{R}^{N+1,1}$:

$$A_\psi(v) = \psi^{-1}v\psi. \quad (54)$$

Obviously, for a vector u with $u^2 = -1$ one has:

$$A_u(v) = u^{-1}vu = 2\langle u, v \rangle u - v. \quad (55)$$

Thus, A_u is, up to sign, the reflection in the (Minkowski) hyperplane $u^\perp$ orthogonal to u ; these reflections generate the genuine Lorentz group. The induced Möbius transformation on $\mathbb{S}^N$ (for which the minus sign is irrelevant) is the inversion of $\mathbb{S}^N$ in the hypersphere $\pi(u^\perp \cap \mathcal{L}^{N+1}) \subset \mathbb{S}^N$; inversions in hyperspheres generate the Möbius group. In particular, if u is orthogonal to $\mathbf{e}_\infty$, then A_u fixes $\mathbf{e}_\infty$, hence leaves $\mathcal{K}$ invariant and induces a Euclidean motion on $\mathbb{R}^N$, namely the reflection in an affine hyperplane; such reflections generate the Euclidean group.

Fact 3 *Möbius transformations of $\mathbb{S}^N$ are in a one-to-one correspondence to elements of the group $\mathcal{H}/\{\pm 1\}$; Euclidean transformations of $\mathbb{R}^N$ are in a one-to-one correspondence to elements of $\mathcal{H}_\infty/\{\pm 1\}$.*

4.3 Orthogonal systems in Möbius geometry

We use elements $\psi \in \mathcal{H}_\infty$ as frames when describing orthogonal systems in the Möbius picture. The vectors $A_\psi(\mathbf{e}_i)$ for $i = 1, \dots, N$ form an orthogonal basis in $T\mathcal{K}$ at the point $A_\psi(\mathbf{e}_0) \in \mathcal{K}$. For an orthogonal system $x^\epsilon : B^\epsilon(r) \mapsto \mathcal{K}$, continuous ($\epsilon = 0$) or discrete ($\epsilon > 0$), with ($m' > 0$) or without ($m' = 0$) transformations, consider its lift to the conic section $\mathcal{K}$:

$$\hat{x}^\epsilon = \lambda \circ x^\epsilon : B^\epsilon(r) \rightarrow \mathcal{K}. \quad (56)$$

4.3.1 Continuous orthogonal systems

Define $\hat{v}_i : B_0(r) \rightarrow T\mathcal{K}$ for $i = 1, \dots, m$ by

$$\partial_i \hat{x} = h_i \hat{v}_i, \quad h_i = |\partial_i x|. \quad (57)$$

The vectors $\hat{v}_i$ are pairwise orthogonal, orthogonal to $\mathbf{e}_\infty$, and can be written as

$$\hat{v}_i = v_i + 2\langle x, v_i \rangle \mathbf{e}_\infty \quad (58)$$

with the vector fields v_i from (43). As an immediate consequence, these vectors satisfy

$$\partial_i \hat{v}_j = \beta_{ji} \hat{v}_i, \quad 1 \leq i \neq j \leq m, \quad (59)$$

with the same rotation coefficients β_{ji} as in (44).

Definition 8 *a) Given a point $\hat{x} \in \mathcal{K}$ and vectors $\hat{v}_i \in T_{\hat{x}}\mathcal{K}$, $1 \leq i \leq m$, we call an element $\psi \in \mathcal{H}_\infty$ suited to $(\hat{x}; \hat{v}_1, \dots, \hat{v}_m)$ if*

$$A_\psi(\mathbf{e}_0) = \hat{x}, \quad (60)$$

$$A_\psi(\mathbf{e}_k) = \hat{v}_k \quad (1 \leq k \leq m), \quad (61)$$

b) A frame $\psi : B_0(r) \mapsto \mathcal{H}_\infty$ is called adapted to the orthogonal system $\hat{x}$, if it is suited to $(\hat{x}; \hat{v}_1, \dots, \hat{v}_m)$ at any point $\xi \in B_0(r)$, and

$$\partial_i A_\psi(\mathbf{e}_k) = \beta_{ki} \hat{v}_i \quad (1 \leq k \leq N, \quad 1 \leq i \leq m). \quad (62)$$

with some scalar functions β_{ki} .

Proposition 3 *For a given continuous orthogonal system $\hat{x} : B_0(r) \mapsto \mathcal{K}$, an adapted frame ψ always exists. It is unique, if $m = N$. If $m < N$, then an adapted frame is uniquely determined by its value at one point, say $\xi = 0$. An adapted frame ψ satisfies the following differential equations.*

- *Static frame equations:*

$$\partial_i \psi = -\mathbf{e}_i \psi s_i \quad (1 \leq i \leq m) \quad (63)$$

with $s_i = (1/2)\partial_i \hat{v}_i$.

- *Moving frame equations*

$$\partial_i \psi = -S_i \mathbf{e}_i \psi \quad (1 \leq i \leq m) \quad (64)$$

with

$$S_i = A_{\mathbf{e}_i\psi}^{-1}(s_i) = \frac{1}{2} \sum_{k \neq i} \beta_{ki} \mathbf{e}_k - h_i \mathbf{e}_\infty, \quad (65)$$

where the functions β_{ij} and h_i constitute a solution to the Lamé system (in the following equations the indices $1 \leq i, j \leq m$ and $1 \leq k \leq N$ are pairwise distinct):

$$\partial_i h_j = h_i \beta_{ij}, \quad (66)$$

$$\partial_i \beta_{kj} = \beta_{ki} \beta_{ij}, \quad (67)$$

$$\partial_i \beta_{ij} + \partial_j \beta_{ji} = - \sum_{k \neq i, j} \beta_{ki} \beta_{kj}. \quad (68)$$

Conversely, if β_{ij} and h_i are solutions to the equations (66)-(68), then the moving frame equations (64) are compatible, and any solution is a frame adapted to an orthogonal system.

Proof. For an orthogonal coordinate system ($m = N$) the adapted frame is uniquely determined at any point by the requirements (60) and (61), while eqs. (62) follow from (61). If $m < N$, extend the m vectors $\hat{v}_i(0)$, $1 \leq i \leq m$, by $N - m$ vectors $\hat{v}_k(0)$, $m < k \leq N$, to a orthonormal basis of $T_{\hat{x}(0)}\mathcal{K}$. There exist unique vector fields $\hat{v}_k : B_0(r) \mapsto T\mathcal{K}$, $m < k \leq N$ with these prescribed values at $\xi = 0$, normalized ($\hat{v}_k^2 = -1$), orthogonal to each other, to $\hat{v}_i$, $1 \leq i \leq m$, to $\hat{x}$ and to $\mathbf{e}_\infty$, and satisfying differential equations

$$\partial_i \hat{v}_k = \beta_{ki} \hat{v}_i, \quad 1 \leq i \leq m, \quad m < k \leq N \quad (69)$$

with some scalar functions β_{ki} . Indeed, orthogonality conditions yield that necessarily $\beta_{ki} = -\langle \partial_i \hat{v}_i, \hat{v}_k \rangle$. Eqs. (69) with these expressions for β_{ki} form a well-defined linear system of first order partial differential equations for $\hat{v}_k$. The compatibility condition for this system, $\partial_j(\partial_i \hat{v}_k) = \partial_j(\partial_i \hat{v}_k)$ ($1 \leq i \neq j \leq m$), is easily verified using (59). The unique frame $\psi : B_0(r) \mapsto \mathcal{H}_\infty$ with $A_\psi(\mathbf{e}_0) = \hat{x}$, $A_\psi(\mathbf{e}_k) = \hat{v}_k$ ($1 \leq k \leq N$) is adapted to $\hat{x}$.

Next, we prove the statements about the moving frame equation. The equations

$$\partial_i A_\psi(\mathbf{e}_0) = h_i A_\psi(\mathbf{e}_i), \quad \partial_i A_\psi(\mathbf{e}_k) = \beta_{ki} A_\psi(\mathbf{e}_i)$$

are equivalent to

$$[\mathbf{e}_0, (\partial_i \psi) \psi^{-1}] = h_i \mathbf{e}_i, \quad [\mathbf{e}_k, (\partial_i \psi) \psi^{-1}] = \beta_{ki} \mathbf{e}_i.$$

Since, by (53), $(\partial_i \psi)^{-1} \psi$ is spanned by bivectors, we come to the representation of this element in the form $(\partial_i \psi)^{-1} \psi = -S_i \mathbf{e}_i$ with S_i as in (65). The

compatibility conditions for the moving frame equations, $\partial_i(\partial_j\psi) = \partial_j(\partial_i\psi)$, are equivalent to

$$\partial_j S_i \mathbf{e}_j + \partial_i S_j \mathbf{e}_i + S_i(\mathbf{e}_i S_j \mathbf{e}_i) + S_j(\mathbf{e}_j S_i \mathbf{e}_j) = 0,$$

which, in turn, are equivalent to the system (66)-(68). Conversely, for a solution $\psi : B_0(r) \mapsto \mathcal{H}_\infty$ of the moving frame equations, define $\hat{x} = A_\psi(\mathbf{e}_0)$ and $\hat{v}_k = A_\psi(\mathbf{e}_k)$. Then $\partial_i \hat{x} = h_i \hat{v}_i$, and orthogonality of $\hat{v}_i$ yields that $\hat{x}$ is an orthogonal system.

We conclude by showing the static frame equation. If ψ is adapted, then (65) in combination with the orthogonality relation of the v_i yields:

$$\begin{aligned} s_i &= A_{\mathbf{e}_i\psi}(S_i) = \frac{1}{2} \sum_{k \neq i} \beta_{ki} A_{\mathbf{e}_i\psi}(\mathbf{e}_k) - h_i A_{\mathbf{e}_i\psi}(\mathbf{e}_\infty) \\ &= -\frac{1}{2} \sum_{k \neq i} \beta_{ki} A_\psi(\mathbf{e}_k) + h_i A_\psi(\mathbf{e}_\infty) = -\frac{1}{2} \sum_{k \neq i} \beta_{ki} \hat{v}_k + h_i \mathbf{e}_\infty = \frac{1}{2} \partial_i \hat{v}_i. \end{aligned}$$

The last equality follows from $\beta_{ki} = -\langle \partial_i \hat{v}_i, \hat{v}_k \rangle$ and the fact that the $\mathbf{e}_\infty$ -component of $\partial_i \hat{v}_i$ is equal to $2h_i$ (the latter follows from (58)). ■

4.3.2 Discrete orthogonal systems

Now consider a discrete orthogonal system $\hat{x} : B^\epsilon(r) \mapsto \mathcal{K}$. Recall that the cases $m' = 0$ and $m' > 0$ are treated simultaneously, so that $\epsilon_i = \epsilon$ for $1 \leq i \leq m$, and $\epsilon_i = 1$ for $m+1 \leq i \leq m+m' = M$. Appropriate discrete analogues of the metric coefficients h_i and vectors $\hat{v}_i$ are

$$h_i = |\delta_i \hat{x}|, \quad \hat{v}_i = h_i^{-1} \delta_i \hat{x}. \quad (70)$$

These are unit vectors, i.e. $\hat{v}_i^2 = -1$, representing the reflections taking $\hat{x}$ to $\tau_i \hat{x}$ in the Möbius picture, cf. Fig. 5:

$$\tau_i \hat{x} = -A_{\hat{v}_i} \hat{x} = \hat{x} + \epsilon_i h_i \hat{v}_i. \quad (71)$$

It is important to note that vectors $\hat{v}_i$ are not mutually orthogonal. Next, we derive equations that are discrete analogues of (59). Since four points $\hat{x}$, $\tau_i \hat{x}$, $\tau_j \hat{x}$ and $\tau_i \tau_j \hat{x}$ are coplanar, the vectors $\tau_j \delta_i \hat{x}$ and $\tau_i \delta_j \hat{x}$ lie in the span of $\delta_i \hat{x}$ and $\delta_j \hat{x}$. The lift λ is affine, therefore the same holds for $\hat{v}_i$'s:

$$\tau_i \hat{v}_j = (n_{ji})^{-1} \cdot (\hat{v}_j + \rho_{ji} \hat{v}_i). \quad (72)$$

Here ρ_{ji} are discrete rotation coefficients, with expected limit behaviour $\rho_{ji} \approx \epsilon_i \epsilon_j \beta_{ji}$ as $\epsilon \rightarrow 0$, and

$$n_{ji}^2 = 1 + \rho_{ji}^2 + 2\rho_{ji} \langle \hat{v}_i, \hat{v}_j \rangle \quad (73)$$




Figure 5: An elementary quadrilateral of a discrete orthogonal system

is the normalizing factor that is expected to converge to 1 as $\epsilon \rightarrow 0$. Circularity implies that the angle between $\delta_i \hat{x}$ and $\delta_j \hat{x}$ and the angle between $\tau_j \delta_i \hat{x}$ and $\tau_i \delta_j \hat{x}$ sum up to π :

$$\langle \tau_j \hat{v}_i, \hat{v}_j \rangle + \langle \hat{v}_i, \tau_i \hat{v}_j \rangle = 0. \quad (74)$$

Under the coplanarity condition (72) the latter condition is equivalent to

$$\hat{v}_j(\tau_j \hat{v}_i) + \hat{v}_i(\tau_i \hat{v}_j) = 0. \quad (75)$$

Inserting (72) and (73) into (74), one finds that either $\langle \hat{v}_i, \hat{v}_j \rangle = 1$, i.e., the circle degenerates to a line (we exclude this from consideration), or

$$\rho_{ij} + \rho_{ji} + 2\langle \hat{v}_i, \hat{v}_j \rangle = 0. \quad (76)$$

From (73) and (76) there follows:

$$n_{ij}^2 = n_{ji}^2 = 1 - \rho_{ij}\rho_{ji}. \quad (77)$$

So, the orthogonality constraint in the discrete case is expressed as the condition (76) on the rotation coefficients; recall that in the continuous case this was a condition (47), resp. (48), on *the derivatives* of the rotation coefficients.

The following definition mimics the static frame equations of the continuous case.

Definition 9 A frame $\psi : B^\epsilon(r) \mapsto \mathcal{H}_\infty$ is called *adapted to a discrete orthogonal system* $\hat{x}$, if

$$A_\psi(\mathbf{e}_0) = \hat{x}, \quad (78)$$

$$\tau_i \psi = -\mathbf{e}_i \psi \hat{v}_i \quad (1 \leq i \leq M). \quad (79)$$

Existence of such a frame is a consequence of the circularity condition. Indeed, the consistency $\tau_j \tau_i \psi = \tau_i \tau_j \psi$ is written as

$$\mathbf{e}_i \mathbf{e}_j \psi \hat{v}_j (\tau_j \hat{v}_i) = \mathbf{e}_j \mathbf{e}_i \psi \hat{v}_i (\tau_i \hat{v}_j).$$

which is equivalent to (75). An adapted frame is uniquely defined by the choice of $\psi(0)$ (and thus is not unique, even if $M = N$). Indeed, eq. (79) together with the definition (70) of $\hat{v}_i$ imply that if eq. (78) holds at one point of $B^\epsilon(r)$, then it holds everywhere.

Proposition 4 *An adapted frame ψ for a discrete orthogonal system $\hat{x}$ satisfies the moving frame equations:*

$$\tau_i \psi = -\Sigma_i \mathbf{e}_i \psi \quad (1 \leq i \leq M) \quad (80)$$

with

$$\Sigma_i = A_{\mathbf{e}_i \psi}^{-1}(\hat{v}_i) = N_i \mathbf{e}_i + \frac{\epsilon_i}{2} \sum_{k \neq i} \beta_{ki} \mathbf{e}_k - \epsilon_i h_i \mathbf{e}_\infty, \quad (81)$$

where

$$N_i^2 = 1 - \frac{\epsilon_i^2}{4} \sum_{k \neq i} \beta_{ki}^2, \quad (82)$$

and the functions β_{ij} and h_i solve the following discrete Lamé system (in the equations below $1 \leq i, j \leq M$, $1 \leq k \leq N$, and $i \neq j \neq k \neq i$):

$$\tau_i h_j = (n_{ji} \epsilon_j)^{-1} \cdot (\epsilon_j h_j + \epsilon_i h_i \rho_{ij}), \quad (83)$$

$$\tau_i \beta_{kj} = (n_{ji} \epsilon_j)^{-1} \cdot (\epsilon_j \beta_{kj} + \epsilon_i \beta_{ki} \rho_{ij}), \quad (84)$$

$$\tau_i \beta_{ij} = (n_{ji} \epsilon_j)^{-1} \cdot (2N_i \rho_{ij} - \epsilon_j \beta_{ij}), \quad (85)$$

$$\rho_{ij} + \rho_{ji} = N_i \beta_{ij} \epsilon_j + N_j \beta_{ji} \epsilon_i - \frac{\epsilon_i \epsilon_j}{2} \sum_{k \neq i, j} \beta_{ki} \beta_{kj}. \quad (86)$$

where the abbreviation $n_{ij} = (1 - \rho_{ij} \rho_{ji})^{1/2}$ is used, and ρ_{ij} are suitable real-valued functions. Conversely, given a solution β_{ij} and h_i of the equations (83)-(86), the system of moving frame equations (80) is consistent, and its solution ψ is an adapted frame of a discrete orthogonal system.

Proof. From the definition $\Sigma_i = A_{\mathbf{e}_i \psi}^{-1}(\hat{v}_i)$ it follows that Σ_i is a vector without $\mathbf{e}_0$ -component, i.e. admits a decomposition of the form (81). Again, from the definition of Σ_i and the moving frame equation (79) we find:

$$A_{\mathbf{e}_i}(\Sigma_j) = \mathbf{e}_i \mathbf{e}_j \psi \hat{v}_j \psi^{-1} \mathbf{e}_j \mathbf{e}_i, \quad \tau_i \Sigma_j = -\mathbf{e}_j \mathbf{e}_i \psi \hat{v}_i (\tau_i \hat{v}_j) \hat{v}_i \psi^{-1} \mathbf{e}_i \mathbf{e}_j.$$

Upon using the identity $\hat{v}_i(\hat{v}_j + \rho_{ji}\hat{v}_i)\hat{v}_i = \hat{v}_j + \rho_{ij}\hat{v}_i$, which follows easily from (76), we can now represent eq. (72) in the following equivalent form:

$$n_{ji}(\tau_i \Sigma_j) = -A_{\mathbf{e}_i}(\Sigma_j) - \rho_{ij}A_{\mathbf{e}_j}(\Sigma_i). \quad (87)$$

This equation is equivalent to eqs. (83)–(85). Next, the consistency $\tau_j(\tau_i \psi) = \tau_i(\tau_j \psi)$ of the moving frame equations (80) is equivalent to

$$(\tau_i \Sigma_j)A_{\mathbf{e}_j}(\Sigma_i) + (\tau_j \Sigma_i)A_{\mathbf{e}_i}(\Sigma_j) = 0. \quad (88)$$

Inserting the expressions for $\tau_i \Sigma_j$ and $\tau_j \Sigma_i$ from (87) into the left side give:

$$\begin{aligned} & A_{\mathbf{e}_i}(\Sigma_j)A_{\mathbf{e}_j}(\Sigma_i) + A_{\mathbf{e}_j}(\Sigma_i)A_{\mathbf{e}_i}(\Sigma_j) + \rho_{ji}(A_{\mathbf{e}_i}(\Sigma_j))^2 + \rho_{ij}(A_{\mathbf{e}_j}(\Sigma_i))^2 \\ &= N_i \beta_{ji} \epsilon_i + N_j \beta_{ij} \epsilon_j - (\epsilon_i \epsilon_j / 2) \sum_{k \neq i, j} \beta_{ki} \beta_{kj} - (\rho_{ij} + \rho_{ji}) = 0, \end{aligned}$$

the last equality being nothing but eq. (86). Conversely, eqs. (83)–(86) imply (87), as well as the compatibility of the frame equations (80). So for an arbitrary initial value $\psi(0) \in \mathcal{H}_\infty$, the frame equations for $\psi : B^\epsilon(r) \mapsto \mathcal{H}_\infty$ can be solved uniquely. Set $\hat{x}^\epsilon = A_\psi(\mathbf{e}_0)$, $\hat{v}_i = A_{\mathbf{e}_i \psi}(\Sigma_i)$. From (87), (88) for the quantities Σ_i there follow (72), (74) for the quantities $\hat{v}_i$. Using the fact that Σ_i has no $\mathbf{e}_0$ -component, one shows easily that there holds also (71). Therefore, $\hat{x}$ is a discrete orthogonal system. ■

4.3.3 Ribaucour transformation of a continuous orthogonal system

To describe a Ribaucour pair $\hat{x}, \hat{x}^+ : B_0(r) \mapsto \mathcal{K}$ of continuous orthogonal systems, one combines the descriptions of two previous subsections. Recall that in the present context $M = m + 1$.

We denote by h_i^+ , $\hat{v}_i^+$, β_{ij}^+ the corresponding objects for $\hat{x}^+$, defined as in (57), (59). Denote by $\hat{v}_M$ unit vectors such that

$$\hat{x}^+ = -A_{\hat{v}_M}(\hat{x}), \quad (89)$$

cf. eq. (71). The defining property (42) of Ribaucour transformations and the normalization of $\hat{v}_M$ imply:

$$\hat{v}_i^+ = -A_{\hat{v}_M}(\hat{v}_i), \quad (90)$$

$$\partial_i \hat{v}_M = \frac{\alpha_i}{2} (\hat{v}_i^+ + \hat{v}_i) = \alpha_i (\hat{v}_i - \langle \hat{v}_M, \hat{v}_i \rangle \hat{v}_M), \quad (91)$$

with m auxiliary functions $\alpha_i : B_0(r) \mapsto \mathbb{R}$.

Definition 10 A pair of frames $\psi, \psi^+ : B_0(r) \mapsto \mathcal{H}_\infty$ is called adapted to the Ribaucour pair $\hat{x}, \hat{x}^+$, if ψ is adapted to $\hat{x}$ and

$$\psi^+ = -\mathbf{e}_M \psi \hat{v}_M. \quad (92)$$

Proposition 5 Let ψ, ψ^+ be a pair of frames adapted to the Ribaucour pair $\hat{x}, \hat{x}^+$. Then the frame ψ^+ is adapted to $\hat{x}^+$, and the following moving frame equations hold for ψ^+ :

$$\partial_i \psi^+ = -S_i^+ \mathbf{e}_i \psi^+ \quad (1 \leq i \leq m), \quad (93)$$

$$\psi^+ = -\Sigma_M \mathbf{e}_M \psi, \quad (94)$$

where

$$S_i^+ = \frac{1}{2} \sum_{k \neq i} \beta_{ki}^+ \mathbf{e}_k - h_i^+ \mathbf{e}_\infty \quad (1 \leq i \leq m), \quad (95)$$

$$\Sigma_M = A_{\mathbf{e}_M \psi}^{-1}(\hat{v}_M) = N_M \mathbf{e}_M + \frac{1}{2} \sum_{k \neq M} \beta_{kM} \mathbf{e}_k - h_M \mathbf{e}_\infty, \quad (96)$$

$$N_M^2 = 1 - \frac{1}{4} \sum_{k \neq M} \beta_{kM}^2, \quad (97)$$

the functions h_i, β_{ij} and h_i^+, β_{ij}^+ solve eqs. (66)-(68), and the following system is satisfied (for $1 \leq i \neq j \leq m$, $1 \leq k \leq N$, and $i \neq k \neq M \neq i$):

$$h_i^+ = h_i + h_M \alpha_i, \quad (98)$$

$$\beta_{ki}^+ = \beta_{ki} + \beta_{kM} \alpha_i, \quad (99)$$

$$\beta_{Mi}^+ = -\beta_{Mi} + 2N_M \alpha_i, \quad (100)$$

$$\partial_i h_M = \frac{1}{2} (h_i + h_i^+) \beta_{iM}, \quad (101)$$

$$\partial_i \beta_{kM} = \frac{1}{2} (\beta_{ki} + \beta_{ki}^+) \beta_{iM}, \quad (102)$$

$$\partial_i \beta_{iM} = -\frac{1}{2} \sum_{k \neq i, M} (\beta_{ki} + \beta_{ki}^+) \beta_{kM} + N_M (\beta_{Mi} - \beta_{Mi}^+), \quad (103)$$

$$\partial_i \alpha_j = \frac{1}{2} \alpha_i (\beta_{ij} + \beta_{ij}^+). \quad (104)$$

Conversely, given solutions h_i, β_{ij} and h_i^+, β_{ij}^+ of the system above with suitable auxiliary functions α_i , the moving frame equations (93), (94) are compatible, and the solution ψ, ψ^+ is a pair of frames adapted to a Ribaucour pair of orthogonal systems.

Proof. First we show that ψ^+ is an adapted frame for x^+ . We have for $1 \leq i \leq m, 1 \leq k \leq N$:

$$\begin{aligned}
A_{\psi^+}(\mathbf{e}_0) &= A_{\hat{v}_M} A_{\mathbf{e}_M \psi}(\mathbf{e}_0) = -A_{\hat{v}_M}(\hat{x}) = \hat{x}^+, \\
A_{\psi^+}(\mathbf{e}_i) &= A_{\hat{v}_M} A_{\mathbf{e}_M \psi}(\mathbf{e}_i) = -A_{\hat{v}_M}(\hat{v}_i) = \hat{v}_i^+, \\
\partial_i A_{\psi^+}(\mathbf{e}_k) &= \partial_i (A_{\hat{v}_M} A_{\mathbf{e}_M \psi}(\mathbf{e}_k)) = \pm \partial_i (A_{\hat{v}_M} A_{\psi}(\mathbf{e}_k)) \\
&= \pm \partial_i (A_{\psi}(\mathbf{e}_k) - 2\langle \hat{v}_M, A_{\psi}(\mathbf{e}_k) \rangle \hat{v}_M) \\
&= \pm (\beta_{ki} - 2\alpha_i \langle \hat{v}_M, A_{\psi}(\mathbf{e}_k) \rangle) (\hat{v}_i - 2\langle \hat{v}_M, \hat{v}_i \rangle \hat{v}_M) \\
&= \pm (\beta_{ki} - 2\alpha_i \langle \hat{v}_M, A_{\psi}(\mathbf{e}_k) \rangle) \hat{v}_i^+ = \beta_{ki}^+ \hat{v}_i^+,
\end{aligned}$$

where the minus sign applies iff $k = M$ (and if $k = 0$, the latter calculation goes through almost literally, with replacing β_{ki}, β_{ki}^+ by h_i, h^+ , respectively). Next, from $\Sigma_M = A_{\mathbf{e}_M \psi}^{-1}(\hat{v}_M)$ there follows:

$$\begin{aligned}
\beta_{kM} &= 2\langle \Sigma_M, \mathbf{e}_k \rangle = 2\langle \hat{v}_M, A_{\mathbf{e}_M \psi}(\mathbf{e}_k) \rangle = -2\langle \hat{v}_M, A_{\psi}(\mathbf{e}_k) \rangle, \\
h_M &= 2\langle \Sigma_M, \mathbf{e}_0 \rangle = 2\langle \hat{v}_M, A_{\mathbf{e}_M \psi}(\mathbf{e}_0) \rangle = -2\langle \hat{v}_M, x \rangle, \\
N_M &= \langle \Sigma_M, \mathbf{e}_M \rangle = \langle \hat{v}_M, A_{\mathbf{e}_M \psi}(\mathbf{e}_M) \rangle = \langle \hat{v}_M, A_{\psi}(\mathbf{e}_M) \rangle.
\end{aligned}$$

This proves eqs. (98)–(100). Further, eqs. (101)–(103) are readily derived by calculating the i -th partial derivative of the respective scalar product. Finally, eq. (104) comes from the consistency condition $\partial_i(\partial_j \hat{v}_M) = \partial_j(\partial_i \hat{v}_M)$.

Conversely, given a solution to the equations (66)–(68) and (98)–(104), the moving frame equations are consistent, thus defining the frames ψ, ψ^+ . It follows from Proposition 3 that both $\hat{x} = A_{\psi}(\mathbf{e}_0)$ and $\hat{x}^+ = A_{\psi^+}(\mathbf{e}_0)$ are orthogonal systems. Furthermore, eq. (94), yields the defining relations (89)–(91) of the Ribaucour pair, with $\hat{v}_M = A_{\mathbf{e}_M \psi}(\Sigma_M)$. ■

5 Goursat problems and approximation for orthogonal systems

It would be tempting to derive the theory of the Lamé system (43)–(47) from its discrete counterpart (83)–(86), treating the latter as a hyperbolic system of the type (11). However, it turns out that in dimensions $M \geq 3$ this is hard to carry out, since one needs to enlarge the set of dependent variables and equations in a cumbersome manner. The way around is based on the following fundamental lemma, which allows one to take care of two-dimensional orthogonal systems (in the coordinate planes) only, and to use then the results for conjugate systems.

Lemma 3 *a) If for a conjugate net x , continuous or discrete, its restriction to each plane $\mathcal{P}_{ij}$, $1 \leq i \neq j \leq M$, is a C-surface, then x is a (continuous or discrete) orthogonal system.*

b) If in a Jonas pair (x, x^+) , continuous or discrete, the net x is an orthogonal system, and the corresponding coordinate curves $x|_{\mathcal{P}_i}$ and $x^+|_{\mathcal{P}_i}$ envelope one-dimensional families of circles, then (x, x^+) is a Ribaucour pair of orthogonal systems.

Proof in the discrete case is based on the Miguel theorem, cf. Sect. 1, and can be found in [CDS]. For the continuous case it is a by-product of the proof of Theorem 6 below. ■

So, suppose that $M = 2$. Then the moving frame equations (80) and the system (83)-(86) take the form ($1 \leq i, j \leq 2$, $1 \leq k \leq N$, $i \neq j \neq k \neq i$):

$$\delta_i \psi = \left(\frac{N_i - 1}{\epsilon_i} - \frac{1}{2} \sum_{k \neq i} \beta_{ki} \mathbf{e}_k \mathbf{e}_i + h_i \mathbf{e}_\infty \mathbf{e}_i \right) \psi, \quad (105)$$

$$\delta_i h_j = \frac{\rho_{ij}}{\epsilon_j n} h_i + \frac{1 - n}{\epsilon_i n} h_j, \quad (106)$$

$$\delta_i \beta_{ij} = \frac{2N_i \rho_{ij}}{\epsilon_i \epsilon_j n} - \frac{1 + n}{\epsilon_i n} \beta_{ij}, \quad (107)$$

$$\delta_i \beta_{kj} = \frac{1 - n}{\epsilon_i n} \beta_{kj} + \frac{\rho_{ij}}{\epsilon_j n} \beta_{ki}, \quad (108)$$

$$\rho_{12} + \rho_{21} = \epsilon_2 N_1 \beta_{12} + \epsilon_1 N_2 \beta_{21} - \epsilon_1 \epsilon_2 \Theta. \quad (109)$$

Here we use the abbreviations

$$n = n_{12} = n_{21} = \sqrt{1 - \rho_{12} \rho_{21}}, \quad \Theta = \frac{1}{2} \sum_{k > 2} \beta_{k1} \beta_{k2}, \quad N_i^2 = 1 - \frac{\epsilon_i^2}{4} \sum_{k \neq i} \beta_{ki}^2. \quad (110)$$

We show how to re-formulate the above system in the hyperbolic form and to pose a Cauchy problem for it.

5.1 Hyperbolic equations and approximation for C-surfaces

First consider the case $m = 2$, $m' = 0$, $\epsilon_1 = \epsilon_2 = \epsilon$, corresponding to C-surfaces. Set

$$\rho_{12} = \epsilon N_1 \beta_{12} - (\epsilon^2/2)(\Theta - \gamma), \quad (111)$$

$$\rho_{21} = \epsilon N_2 \beta_{21} - (\epsilon^2/2)(\Theta + \gamma), \quad (112)$$

with a suitable function $\gamma : \mathcal{P}_{12}^\epsilon \mapsto \mathbb{R}$. It can be said that the auxiliary function γ splits the constraint, making the Lamé system hyperbolic. This splitting plays a crucial role for our approximation results. In the smooth limit $2\gamma = \partial_1\beta_{12} - \partial_2\beta_{21}$ (cf. eqs. (114), (115) below).

As a Banach space for the system take

$$\mathcal{X} = \hat{\mathcal{H}}\{\psi\} \times \mathbb{R}^{2(N-1)}\{\beta\} \times \mathbb{R}^2\{h_1, h_2\} \times \mathbb{R}\{\gamma\},$$

where $\hat{\mathcal{H}}$ is the space of an (arbitrary) matrix representation of $Spin(N+1, 1)$ (note that if $\psi \in \mathcal{H}_\infty \subset \hat{\mathcal{H}}$ at some point, then eq. (105) guarantees that this is the case everywhere), β denotes the collection of β_{kj} with $k = 1, 2$, $1 \leq j \leq N$, $k \neq j$. Obviously,

$$N_1 = 1 + O(\epsilon^2), \quad N_2 = 1 + O(\epsilon^2), \quad n = 1 + O(\epsilon^2), \quad (113)$$

where the constants in O -symbols are uniform on compact subsets of $\mathcal{X}$.

Both directions $i = 1, 2$ are assumed to be evolution directions for ψ ; for h_i and β_{ki} there is one evolution direction $j \neq i$, while for the additional function γ both directions $i = 1, 2$ are static.

Goursat Problem 3 (for discrete C-Surfaces). *Given $\Psi^\epsilon \in \mathcal{H}_\infty$, functions $H_i^\epsilon : \mathcal{P}_i^\epsilon \mapsto \mathbb{R}$ and $B_{ki}^\epsilon : \mathcal{P}_i^\epsilon \mapsto \mathbb{R}$ for $i = 1, 2$, $1 \leq k \leq N$, $k \neq i$, and $\Gamma^\epsilon : \mathcal{P}_{12}^\epsilon \mapsto \mathbb{R}$, find a solution to the equations (105)-(108) with (111), (112), and with the Goursat data*

$$\psi(0) = \Psi^\epsilon, \quad h_i|_{\mathcal{P}_i^\epsilon} = H_i^\epsilon, \quad \beta_{ki}|_{\mathcal{P}_i^\epsilon} = B_{ki}^\epsilon, \quad \gamma|_{\mathcal{P}_{12}^\epsilon} = \Gamma^\epsilon.$$

Lemma 4 *The hyperbolic system (105)–(108) with (111), (112) is consistent. Goursat problem 3 for discrete C-surfaces satisfies condition (F). The limiting system for $\epsilon = 0$ consists of equations (64) with (65), (66), (67) for pairwise distinct indices $1 \leq i, j \leq 2$, $1 \leq k \leq N$, and*

$$\partial_1\beta_{12} = -(1/2) \sum_{k>2} \beta_{k1}\beta_{k2} + \gamma, \quad (114)$$

$$\partial_2\beta_{21} = -(1/2) \sum_{k>2} \beta_{k1}\beta_{k2} - \gamma, \quad (115)$$

which describe continuous C-surfaces.

Proof. Consistency is easy to see: the only condition to be checked is $\delta_1\delta_2\psi = \delta_2\delta_1\psi$, but this has already been shown in Proposition 4. Also other statements are obvious from (113). Notice that hyperbolic equations (114), (115) come to replace the non-hyperbolic orthogonality constraint (68). ■

Now we discuss the way to prescribe the data in the Goursat problem 3 in order to get an approximation of a given smooth C-surface. Unlike in the case of general conjugate nets, it is not possible to prescribe the discrete curves $X_i^\epsilon = x^\epsilon|_{\mathcal{P}_i^\epsilon}$ coinciding with their continuous counterparts $X_i = x|_{\mathcal{P}_i}$ at the lattice points. However, it is still possible to achieve that X_i^ϵ are completely determined by X_i .

Given a curve $\hat{X}_i : \mathcal{P}_i \mapsto \mathcal{K}$, one has the corresponding tangential vector field $\partial_i \hat{X}_i$, and therefore the function $h_i = |\partial_i \hat{X}_i|$ and the field of unit vectors $\hat{v}_i = h_i^{-1} \partial_i \hat{X}_i$. Take $\Psi \in \mathcal{H}_\infty$ suited to $(\hat{X}_i(0); \hat{v}_i(0))$. Then by Proposition 3, there is a unique frame $\psi : \mathcal{P}_i \rightarrow \mathcal{H}_\infty$ adapted to the curve $\hat{X}_i$ with $\psi(0) = \Psi$; this frame is defined as the unique solution of the differential equation $\partial_i \psi = -S_i \mathbf{e}_i \psi$ with S_i given by eq. (65). According to the proof of Proposition 3, the latter equation is equivalent to the system of equations

$$\partial_i \hat{v}_k = \beta_{ki} \hat{v}_i, \quad \beta_{ki} = -\langle \hat{v}_k, \partial_i \hat{v}_i \rangle, \quad k \neq i.$$

So, in order to determine the rotation coefficients $\beta_{ki} : \mathcal{P}_i \mapsto \mathbb{R}$ for $1 \leq k \leq N$, $k \neq i$, one has to solve the latter system of ordinary differential equations with the initial data $\hat{v}_k(0) = A_\Psi(\mathbf{e}_k)$. Thus, we produced the functions h_i and β_{ki} , or, what is equivalent, the Clifford elements

$$S_i = \frac{1}{2} \sum_{k \neq i} \beta_{ki} \mathbf{e}_k - h_i \mathbf{e}_\infty$$

for a given curve $\hat{X}_i : \mathcal{P}_i \mapsto \mathcal{K}$. We say that h_i, β_{ki} are *read off* the curve $\hat{X}_i$.

Definition 11 *The canonical discretization of the curve $\hat{X}_i : \mathcal{P}_i \mapsto \mathcal{K}$ with respect to the initial frame $\Psi \in \mathcal{H}_\infty$ suited to $(\hat{X}_i(0); \hat{v}_i(0))$ is the function $\hat{X}_i^\epsilon = A_{\psi^\epsilon}(\mathbf{e}_0) : \mathcal{P}_i^\epsilon \mapsto \mathcal{K}$, where $\psi^\epsilon : \mathcal{P}_i^\epsilon \mapsto \mathcal{H}_\infty$ is the solution of the discrete moving frame equation $\tau_i \psi^\epsilon = -\Sigma_i \mathbf{e}_i \psi^\epsilon$ with*

$$\Sigma_i = \left(1 - \frac{\epsilon^2}{4} \sum_{k \neq i} \beta_{ki}^2 \right)^{1/2} \mathbf{e}_i + \frac{\epsilon}{2} \sum_{k \neq i} \beta_{ki} \mathbf{e}_k - \epsilon h_i \mathbf{e}_\infty$$

and the initial condition $\psi^\epsilon(0) = \Psi$.

In other words, for the canonical discretization the data h_i and β_{ki} are read off the continuous curve. Obviously, ψ^ϵ is an adapted frame for the discrete curve X_i^ϵ . As $\epsilon \rightarrow 0$, the canonical discretization X_i^ϵ converges to X_i with the rate $O(\epsilon)$ in C^∞ .

Proposition 6 (approximation for C-surfaces). *Let $m = M = 2$, and let there be given:*

- two smooth curves $X_i : \mathcal{P}_i \mapsto \mathbb{R}^N$ ($i = 1, 2$), intersecting orthogonally at $X = X_1(0) = X_2(0)$,
- a smooth function $\Gamma : \mathcal{P}_{12} \mapsto \mathbb{R}$.

Assume $\Psi \in \mathcal{H}_\infty$ is suited for $(\hat{X}; \hat{v}_1(0), \hat{v}_2(0))$. Then, for some $r > 0$:

1. There exists a unique C -surface $x : B_0(r) \mapsto \mathbb{R}^N$ that coincides with X_i on $\mathcal{P}_i \cap B_0(r)$ and satisfies

$$\partial_1 \beta_{12} - \partial_2 \beta_{21} = 2\Gamma.$$

2. Consider the family of discrete C -surfaces $\{x^\epsilon : B_0^\epsilon(r) \rightarrow \mathbb{R}^N\}_{0 < \epsilon < \epsilon_1}$ defined as the solutions to the Goursat problem 3 with the data

$$\Psi^\epsilon = \Psi, \quad H_i^\epsilon = H_i|_{\mathcal{P}_i^\epsilon}, \quad B_{ki}^\epsilon = B_{ki}|_{\mathcal{P}_i^\epsilon}, \quad \Gamma^\epsilon = \Gamma|_{\mathcal{P}_{12}^\epsilon},$$

where the functions H_i and B_{ki} are read off the smooth curves X_i , so that $x^\epsilon|_{\mathcal{P}_i^\epsilon}$ are the canonical ϵ -discretizations of X_i . This family of discrete C -surfaces $O(\epsilon)$ -converges in C^∞ to x .

Proof follows from Lemma 4 and Theorem 2. ■




Figure 6: Two curves determine a discrete C -surface via Proposition 6.

5.2 Goursat problem and approximation for an orthogonal system

Theorem 6 (approximation of an orthogonal system). Let $m(m-1)/2$ smooth C -surfaces $\mathcal{S}_{ij} : \mathcal{P}_{ij} \mapsto \mathbb{R}^N$ be given, labelled by $1 \leq i \neq j \leq m$.

m with $\mathcal{S}_{ij} = \mathcal{S}_{ji}$. Assume that for a given $1 \leq i \leq m$ all surfaces $\mathcal{S}_{ij}$ intersect along the curvature lines $X_i = \mathcal{S}_{ij} \upharpoonright_{\mathcal{P}_i}$. Set $h_i = |\partial_i X_i|$, assume that $h_i(0) \neq 0$, and set further $v_i = h_i^{-1} \partial_i X_i$. All curves X_i intersect at one point $X = X_1(0) = \dots = X_m(0)$ orthogonally. Let $\Psi \in \mathcal{H}_\infty(N)$ be suited to $(\hat{X}, \hat{v}_1(0), \dots, \hat{v}_m(0))$. Construct the discrete C-surfaces $\mathcal{S}_{ij}^\epsilon : \mathcal{P}_{ij}^\epsilon \mapsto \mathbb{R}^N$ according to Proposition 6, with the functions $\Gamma_{ij} = (\partial_i \beta_{ij} - \partial_j \beta_{ji})/2$ for $1 \leq i < j \leq m$. Then for some $r > 0$:

1. There exists a unique orthogonal system $x : B_0(r) \mapsto \mathbb{R}^N$, coinciding with $\mathcal{S}_{ij}$ on $\mathcal{P}_{ij} \cap B_0(r)$.
2. There exists a unique family of discrete orthogonal systems $\{x^\epsilon : B_0^\epsilon(r) \mapsto \mathbb{R}^N\}_{0 < \epsilon < \epsilon_0}$ coinciding with $\mathcal{S}_{ij}^\epsilon$ on $\mathcal{P}_{ij}^\epsilon \cap B_0^\epsilon(r)$. The family x^ϵ converges to x with the rate $O(\epsilon)$ in C^∞ .

Proof. C-surfaces are special two-dimensional conjugate nets, therefore the surfaces $\mathcal{S}_{ij}$ can be supplied with the respective coefficients $C_{ij}, C_{ji} : \mathcal{P}_{ij} \mapsto \mathbb{R}$. Now Theorem 3 can be applied. It yields the existence of a unique conjugate net x which has X_i as image of the i -th coordinate axes and C_{ij}, C_{ji} as coefficients on the respective $\mathcal{P}_{ij}$.

Similarly, discrete C-surfaces are special discrete two-dimensional conjugate nets, so for each $\mathcal{S}_{ij}^\epsilon$, the coefficients $C_{ij}^\epsilon, C_{ji}^\epsilon$ are defined, and they converge in C^∞ to the coefficients C_{ij}, C_{ji} of the respective $\mathcal{S}_{ij}$. Note that for any $1 \leq i \leq m$ the discrete surfaces $\mathcal{S}_{ij}^\epsilon$ intersect along the discrete curve $X_i^\epsilon = \mathcal{S}_{ij}^\epsilon \upharpoonright_{\mathcal{P}_i}$, which is the canonical discretization of the curve X_i . The conclusion 2 of Theorem 3 is still valid if the Goursat data for c_{ij}^ϵ in (32) are taken as C_{ij}^ϵ instead of C_{ij} , i.e. read off $\mathcal{S}_{ij}^\epsilon$ rather than $\mathcal{S}_{ij}$. Thus, Theorem 3 delivers a family of discrete conjugate nets $\{x^\epsilon\}$ that is $O(\epsilon)$ -convergent to x . Since the restrictions of x^ϵ to the coordinate planes $\mathcal{P}_{ij}^\epsilon$ are discrete C-surfaces, the discrete variant of Lemma 3 implies that the nets x^ϵ are actually discrete orthogonal systems. It remains to show that the limiting net x is with necessity an orthogonal one. But this follows immediately from the fact of C^∞ -convergence and eq. (75): indeed, we can conclude that for the net x everywhere holds

$$\hat{v}_j \hat{v}_i + \hat{v}_i \hat{v}_j = 0 \quad \Leftrightarrow \quad \langle \hat{v}_i, \hat{v}_j \rangle = 0,$$

that is, $\partial_i x \cdot \partial_j x = 0$. ■

This result immediately implies Theorem 1 from Introduction: to bring its assumptions into the form of Theorem 6 above, one needs only to introduce a curvature line parametrization on surfaces $\mathcal{F}_i$ such that the curves X_i are parametrized by arc-length.

5.3 Goursat problems and approximation for Ribaucour transformations

Now we consider again a two-dimensional ($M = 2$) discrete orthogonal system described by eqs. (105)–(108), but perform a different continuous limit, leading to two curves enveloping a circle congruence (Fig. 7). In other words,




Figure 7: A pair of curves enveloping a circle congruence

we set $m = m' = 1$, $\epsilon_1 = \epsilon$, $\epsilon_2 = 1$. Recall that in this case

$$B^\epsilon(r) = B_0^\epsilon(r) \times \{0, 1\}, \quad \text{where} \quad B_0^\epsilon(r) = \mathcal{P}_1^\epsilon \cap [0, r] = (\epsilon\mathbb{Z}) \cap [0, r],$$

so that only two lines in $\mathcal{P}_{12}^\epsilon$ are considered. Therefore, a function on $\mathcal{P}_{12}^\epsilon$ is conveniently considered as a pair of functions on $\mathcal{P}_1^\epsilon$, or even as just one function on $\mathcal{P}_1^\epsilon$, if one is interested in $B_0^\epsilon(r) \times \{0\}$ only (this will be the case for the function α below). We denote the shift τ_2 by the superscript “+”. Set

$$\rho_{21} = \epsilon\alpha, \quad \rho_{12} = N_1\beta_{12} + \epsilon(N_2\beta_{21} - \Theta - \alpha); \quad (116)$$

with a suitable function $\alpha : \mathcal{P}_1^\epsilon \mapsto \mathbb{R}$.

As the Banach space where the system lives we choose, exactly as before,

$$\mathcal{X} = \hat{\mathcal{H}}\{\psi\} \times \mathbb{R}^{2(N-1)}\{\beta\} \times \mathbb{R}^2\{h_1, h_2\} \times \mathbb{R}\{\alpha\}.$$

In the present case we have:

$$N_1^2 = 1 + O(\epsilon^2), \quad N_2^2 = 1 - \frac{1}{4} \sum_{k \neq 2} \beta_{k2}^2, \quad n = 1 - \frac{\epsilon}{2} \alpha \beta_{12} + O(\epsilon^2), \quad (117)$$

where the constants in O -symbols are uniform on compact subsets of $\mathcal{X}$. However, the system itself is now well defined not on all of $\mathcal{X}$ but rather on the subset where $N_2^2 > 0$:

$$\mathbb{D} = \left\{ (\psi, \beta, h_1, h_2, \alpha) : \sum_{k \neq 2} \beta_{k2}^2 < 4 \right\} \subset \mathcal{X}. \quad (118)$$

Goursat Problem 4 (for a pair of curves enveloping a circle congruence). Given $\Psi^\epsilon \in \mathcal{H}_\infty$, functions $H_1^\epsilon : \mathcal{P}_1^\epsilon \mapsto \mathbb{R}$ and $B_{k1}^\epsilon : \mathcal{P}_1^\epsilon \mapsto \mathbb{R}$ for $2 \leq k \leq N$, and $A^\epsilon : \mathcal{P}_1^\epsilon \mapsto \mathbb{R}$, as well as the real numbers H_2^ϵ and B_{k2}^ϵ for $1 \leq k \leq N$, $k \neq 2$, find a solution to the equations (105)-(108) with (116) on $B^\epsilon(r)$ with the Goursat data

$$\begin{aligned} \psi(0) &= \Psi^\epsilon, \quad h_1|_{\mathcal{P}_1^\epsilon} = H_1^\epsilon, \quad \beta_{k1}|_{\mathcal{P}_1^\epsilon} = B_{k1}^\epsilon, \quad \alpha|_{\mathcal{P}_1^\epsilon} = A^\epsilon, \\ h_2(0) &= H_2^\epsilon, \quad \beta_{k2}(0) = B_{k2}^\epsilon. \end{aligned}$$

Lemma 5 *The hyperbolic system (105)-(108) with (116) is consistent. Goursat problem 4 satisfies condition (F) on the subset $\mathbb{D}$. The limiting system for $\epsilon = 0$ consists of equations (64) with (65) for $i = 1$, (80) with (81) for $i = 2$, and*

$$\partial_1 h_2 = \beta_{12}(h_1 + \alpha h_2/2), \quad (119)$$

$$\partial_1 \beta_{k2} = \beta_{12}(\beta_{k1} + \alpha \beta_{k2}/2), \quad k > 2, \quad (120)$$

$$\partial_1 \beta_{12} = 2(N_2 \beta_{21} - \Theta - \alpha) + \alpha \beta_{12}^2/2, \quad (121)$$

$$h_1^+ - h_1 = \alpha h_2, \quad (122)$$

$$\beta_{k1}^+ - \beta_{k1} = \alpha \beta_{k2}, \quad k > 2, \quad (123)$$

$$\beta_{21}^+ - \beta_{21} = 2(N_2 \alpha - \beta_{21}), \quad (124)$$

and describes a pair of continuous curves enveloping a circle congruence.

Proof. Consistency is shown exactly as in Lemma 4, the limiting system is calculated directly from (105)-(108) using (117). One sees that the system of the lemma coincides with (98)-(103) for $M = 2$. ■

Proposition 7 (approximation of a pair of curves enveloping a circle congruence). *Let $m = 1$, $M = 2$, and let there be given:*

- a smooth curve $X : \mathcal{P}_1 \mapsto \mathbb{R}^N$,
- a smooth function $A : \mathcal{P}_1 \mapsto \mathbb{R}$,
- a point $X^+(0) \in \mathbb{R}^N$.

Set $H_2(0) = |X^+(0) - X(0)|$, $\hat{v}_2(0) = H_2(0)^{-1}(X^+(0) - X(0))$. Assume that $\Psi \in \mathcal{H}_\infty$ is suited for $(\hat{X}_1(0); \hat{v}_1(0))$, and set $\hat{v}_k(0) = A_\Psi(\mathbf{e}_k)$ for $k \neq 2$. Then, for some $r > 0$:

1. *There exists a unique curve $X^+ : \mathcal{P}_1 \cap B_0(r) \mapsto \mathbb{R}^N$ through the point $X^+(0)$ such that the pair (X, X^+) envelopes a circle congruence, and*

$$|\partial_1 X^+| - |\partial_1 X| = A \cdot |X^+ - X|. \quad (125)$$

2. Consider the family of pairs of discrete curves $(X^\epsilon, (X^\epsilon)^+)$ defined as the solutions to the Goursat problem 4 with the data

$$\begin{aligned}\Psi^\epsilon &= \Psi, \quad H_1^\epsilon = H_1|_{\mathcal{P}_1^\epsilon}, \quad B_{k1}^\epsilon = B_{k1}|_{\mathcal{P}_1^\epsilon}, \quad A^\epsilon = A|_{\mathcal{P}_1^\epsilon}, \\ H_2^\epsilon &= H_2(0), \quad B_{k2}^\epsilon = -2\langle \hat{v}_2(0), \hat{v}_k(0) \rangle.\end{aligned}$$

where the functions H_1 and B_{k1} are read off the smooth curve X , so that X^ϵ is the canonical ϵ -discretization of X . These discrete curves $O(\epsilon)$ -converge in C^∞ to (X, X^+) .

Proof follows from Lemma 5 and Theorem 2. ■

Theorem 7 (approximation of a Ribaucour pair). *Let, in addition to the data of Theorem 6, there be given m curves $X_i^+ : \mathcal{P}_i \mapsto \mathbb{R}^N$ with a common intersection point $X^+ = X_1^+(0) = \dots = X_m^+(0)$, and such that each pair (X_i, X_i^+) envelopes a circle congruence. In addition to the discrete surfaces $\mathcal{S}_{ij}^\epsilon$ from Theorem 6, construct discrete curves $(X_i^\epsilon)^+$ according to Proposition 7, with the functions $A_i = (|\partial_i X_i^+| - |\partial_i X_i|)/|X_i^+ - X_i|$. Then for some $r > 0$:*

1. *There exists a unique Ribaucour pair of orthogonal systems $x, x^+ : B_0(r) \mapsto \mathbb{R}^N$ such that x coincides with $\mathcal{S}_{ij}$ on $\mathcal{P}_{ij} \cap B_0(r)$, and x^+ coincides with X_i^+ on $\mathcal{P}_i \cap B_0(r)$.*
2. *There exists a unique family of Ribaucour pairs of discrete orthogonal systems $\{x^\epsilon : B^\epsilon(r) \mapsto \mathbb{R}^N\}_{0 < \epsilon < \epsilon_0}$ coinciding with $\mathcal{S}_{ij}^\epsilon$ on $(\mathcal{P}_{ij} \times \{0\}) \cap B(r)$ and coinciding with $(X_i^\epsilon)^+$ on $(\mathcal{P}_i^\epsilon \times \{1\}) \cap B(r)$. The family x^ϵ converges to the pairs x, x^+ with the rate $O(\epsilon)$ in C^∞ .*

Proof is similar to that of Theorem 6, with the only change in the concluding argument: for the limiting Jonas pair x, x^+ of orthogonal nets we derive from eq. (75):

$$\hat{v}_M \hat{v}_i^+ + \hat{v}_i \hat{v}_M = 0 \quad \Rightarrow \quad \langle \hat{v}_i + \hat{v}_i^+, \hat{v}_M \rangle = 0,$$

which is the defining property of the Ribaucour pair. ■

Letting not one but two or three directions of the orthogonal system remain discrete in the continuous limit (so that $m' = 2$ or $m' = 3$), one arrives at the following statement.

Theorem 8 (permutability of Ribaucour transformations).

1. Given an m -dimensional orthogonal system $x(\cdot, 0, 0) : B_0(r) \mapsto \mathbb{R}^N$ and its two Ribaucour transformations $x(\cdot, 1, 0) : B_0(r) \mapsto \mathbb{R}^N$ and $x(\cdot, 0, 1) : B_0(r) \mapsto \mathbb{R}^N$, there exists a one-parameter family of orthogonal systems $x(\cdot, 1, 1) : B_0(r) \mapsto \mathbb{R}^N$ that are Ribaucour transformations of both $x(\cdot, 1, 0)$ and $x(\cdot, 0, 1)$. Corresponding points of the four conjugate nets are concircular.

2. Given three Ribaucour transformations

$$x(\cdot, 1, 0, 0), x(\cdot, 0, 1, 0), x(\cdot, 0, 0, 1) : B_0(r) \mapsto \mathbb{R}^N$$

of a given m -dimensional orthogonal system $x(\cdot, 0, 0, 0) : B_0(r) \mapsto \mathbb{R}^N$, as well as three further orthogonal systems

$$x(\cdot, 1, 1, 0), x(\cdot, 0, 1, 1), x(\cdot, 1, 0, 1) : B_0(r) \mapsto \mathbb{R}^N$$

such that $x(\cdot, 1, 1, 0)$ is a Ribaucour transformation of both $x(\cdot, 1, 0, 0)$ and $x(\cdot, 0, 1, 0)$ etc., there exists generically a unique orthogonal system

$$x(\cdot, 1, 1, 1) : B_0(r) \mapsto \mathbb{R}^N$$

which is a Ribaucour transformation of all three $x(\cdot, 1, 1, 0), x(\cdot, 0, 1, 1)$ and $x(\cdot, 1, 0, 1)$.

5.4 Example: elliptic coordinates

The simplest nontrivial example, to which the above theory can be applied, is the approximation of two-dimensional conformal maps by circular patterns. Starting with a conformal map $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, i.e. $M = N = 2$,

$$h = |\partial_1 F| = |\partial_2 F| \quad \text{and} \quad \partial_1 F \cdot \partial_2 F = 0,$$

one calculates the metric and rotation coefficients according to equations (43) and (45), and the quantity γ according to (114):

$$h_1 = h_2 = h, \quad \beta_{12} = \frac{\partial_1 h}{h}, \quad \beta_{21} = \frac{\partial_2 h}{h}, \quad \gamma = \partial_1 \beta_{12} = -\partial_2 \beta_{21}$$

To construct a discrete approximation of F , solve the Goursat problem 3 for $\epsilon > 0$ with the data H_i^ϵ , B_{ij}^ϵ and Γ^ϵ that are close to the values of the respective functions h_i , β_{ij} and γ at the corresponding lattice sites. A good choice is, for example, to prescribe

$$\begin{aligned} H_1^\epsilon(\xi_1) &= h(\xi_1 + \epsilon/2, 0), & H_2^\epsilon(\xi_2) &= h(0, \xi_2 + \epsilon/2), \\ B_{21}^\epsilon(\xi_1) &= \beta_{21}(\xi_1 + \epsilon/2, 0), & B_{12}^\epsilon(\xi_2) &= \beta_{12}(0, \xi_2 + \epsilon/2), \\ \Gamma^\epsilon(\xi_1, \xi_2) &= \gamma(\xi_1 + \epsilon/2, \xi_2 + \epsilon/2), \end{aligned}$$




Figure 8: Approximation of elliptic coordinates, $\epsilon = \pi/10$ and $\epsilon = \pi/20$.

for $\xi_1, \xi_2 = i\epsilon$, $0 \leq i \leq R$. If $\psi^\epsilon : B^\epsilon(\epsilon R) \rightarrow \mathcal{H}_\infty$ is the frame of the respective solution to the system (105)-(108), then the function

$$F^\epsilon : B^\epsilon(R\epsilon) \mapsto \mathbb{R}^2, \quad F^\epsilon(\xi_1, \xi_2) = \pi(\psi(\xi_1, \xi_2)^{-1} \mathbf{e}_0 \psi(\xi_1, \xi_2))$$

is a two-dimensional discrete orthogonal system, i.e., the points $F^\epsilon(\xi_1, \xi_2)$, $F^\epsilon(\xi_1 + \epsilon, \xi_2)$, $F^\epsilon(\xi_1, \xi_2 + \epsilon)$ and $F^\epsilon(\xi_1 + \epsilon, \xi_2 + \epsilon)$ lie on a common circle $C^\epsilon(\xi_1, \xi_2)$ in $\mathbb{R}^2$, and $F^\epsilon(\xi_1, \xi_2) = F(\xi_1, \xi_2) + O(\epsilon)$.

This is illustrated with the planar elliptic coordinate system:

$$F(\xi_1, \xi_2) = (\cosh(\xi_1) \cos(\xi_2), \sinh(\xi_1) \sin(\xi_2)),$$

whose coordinate lines are ellipses and hyperbolas. One finds

$$\begin{aligned} h &= (\sinh^2(\xi_1) + \sin^2(\xi_2))^{1/2}, \\ \beta_{12} &= \sinh(2\xi_1)/(2h^2), \quad \beta_{21} = \sin(2\xi_2)/(2h^2), \\ \gamma &= (1 - \cosh(2\xi_1) \cos(2\xi_2))/(4h^4). \end{aligned}$$

Results are displayed in Fig. 8. Their left sides show the original coordinate lines of F , and on the right sides the circles $C^\epsilon(\xi_1, \xi_2)$ are drawn; each intersection point of two coordinate lines on the left corresponds to an intersection point of four circles on the right. There are small defects (the circles do not close up) on the very right of the pictures since, in contrast to F , the discrete maps F^ϵ are not periodic with respect to ξ_2 .

References

- [AKV] A.A. Akhmetshin, I.M. Krichever, Y.S. Volvovski, *Discrete analogs of the Darboux-Egoroff metrics*, Proc. Steklov Inst. Math., **2** (225) (1999), 16–39.
- [Be] M. Berger, *Geometry I*, Berlin: Springer, 1987.
- [Bi1] L. Bianchi, *Le trasformazioni di Ribaucour dei sistemi n -pli ortogonali e il teorema generale di permutabilità*, Annali di Mat., **27** (3) (1918), 183–257.
- [Bi2] L. Bianchi, *Lezioni di Geometria Differenziale*, Bologna: Zanichelli, 1924.
- [Bo] A.I. Bobenko, *Discrete conformal maps and surfaces*, In: *Symmetries and Integrability of Difference Equations, Proc. SIDE II Conference, Canterbury, July 1-5, 1996*, Eds. P.A. Clarkson, F.W. Nijhoff, Cambridge Univ. Press, 1999, pp. 97–108.
- [BH] A. Bobenko, U. Hertrich-Jeromin, *Orthogonal nets and Clifford algebras*, Tohoku Math. Publ., **20** (2001), 7–22.
- [BMS] A.I. Bobenko, D. Matthes, Yu.B. Suris, *Nonlinear hyperbolic equations in surface theory: integrable discretizations and approximation results*, math.NA/0208042.
- [BP] A.I. Bobenko, U. Pinkall, *Discretization of Surfaces and Integrable Systems*, In: *Discrete Integrable Geometry and Physics*, Eds. A.I. Bobenko, R. Seiler, Oxford: Clarendon Press, 1999, 3–58.
- [CDS] J. Cieřliński, A. Doliwa, P.M. Santini, *The integrable discrete analogues of orthogonal coordinate system are multi-dimensional circular lattices*, Phys. Lett. A, **235** (1997), 480–488.

- [Da] G. Darboux, *Leçons sur les Systèmes Orthogonaux et les Coordonnées Curviligne*, Paris: Gauthier-Villars, 1910.
- [DMS] A. Doliwa, S.V. Manakov, P.M. Santini, $\bar{\partial}$ reductions of the multi-dimensional quadrilateral lattice: the multidimensional circular lattice, *Comm. Math. Phys.*, **196** (1998), 1–18.
- [DS1] A. Doliwa, P.M. Santini, *Multidimensional quadrilateral lattices are integrable*, *Phys. Lett. A*, **233** (1997), 265–372.
- [DS2] A. Doliwa, P.M. Santini, *Integrable discrete geometry: the quadrilateral lattice, its transformations and reductions*, In: *Symmetries and Integrability of Difference Equations, Proc. SIDE III Conference*, Eds. D. Levi, O. Ragnisco, Providence: AMS, 2000, 101–119.
- [Du] B. Dubrovin, *Integrable systems in topological field theory*, *Nucl. Phys. B* **379** (1992), 627–689.
- [GT] E.I. Ganzha, S.P. Tsarev, *An algebraic superposition formula and the completeness of Bäcklund transformations of $(2 + 1)$ -dimensional integrable systems*, *Russ. Math. Surv.*, **51** (1996), 1200–1202.
- [H] U. Hertrich-Jeromin, *Introduction to Möbius Differential Geometry*, London Math. Soc. Lect. Notes Ser., Cambridge University Press, to appear.
- [KS] B.G. Konopelchenko, W.K. Schief, *Three-dimensional integrable lattices in Euclidean spaces: conjugacy and orthogonality*, *Proc. R. Soc. London A*, **454** (1998), 3075–3104.
- [K] I.M. Krichever, *Algebraic-geometric n -orthogonal curvilinear coordinate systems and the solution of associativity equations*, *Funct. Anal. Appl.*, **31** (1997), 25–39.
- [MPS] R.R. Martin, J. de Pont, T.J. Sharrock, *Cyclide surfaces in computer aided design*, In: *The mathematics of surfaces*, Ed. J.A. Gregory, Oxford: Clarendon Press, 1986, pp.253–268.
- [N] A.W. Nutbourne, *The solution of frame matching equation*, In: *The mathematics of surfaces*, Ed. J.A. Gregory, Oxford: Clarendon Press, 1986, pp.233–252.
- [Z] V.I. Zakharov, *Description of the n -orthogonal curvilinear coordinate systems and Hamiltonian integrable systems of hydrodynamic type, I. Integration of the Lamé equations*, *Duke Math. J.*, **94** (1998), 103–139.